

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Karl Reinhard Papers/Publications

Natural Resources, School of

---

2002

# Multidisciplinary Coprolite Analysis

Karl Reinhard

*University of Nebraska-Lincoln, kreinhard1@mac.com*

Dennis R. Danielson

*U.S. Army Central ID Lab, Hickman AFB, Hawaii*

Mark Daniels

Sérgio Augusto de Miranda Chaves

*FIOCRUZ, smiranda@ensp.fiocruz.br*

Follow this and additional works at: <http://digitalcommons.unl.edu/natresreinhard>



Part of the [Archaeological Anthropology Commons](#), [Ecology and Evolutionary Biology Commons](#), [Environmental Public Health Commons](#), [Other Public Health Commons](#), and the [Parasitology Commons](#)

---

Reinhard, Karl; Danielson, Dennis R.; Daniels, Mark; and Chaves, Sérgio Augusto de Miranda, "Multidisciplinary Coprolite Analysis" (2002). *Karl Reinhard Papers/Publications*. 23.

<http://digitalcommons.unl.edu/natresreinhard/23>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Karl Reinhard Papers/Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## BIGHORN CAVE

Test Excavation of a Stratified Dry Shelter  
Mohave County, Arizona

edited by

Phil R. Geib  
Donald R. Keller

with contributions by

Mary Butterwick  
Mark Daniels  
Dennis R. Danielson  
Phil R. Geib  
Dawn Greenwald  
Mark Hovezak  
Richard H. Hevly  
Chris Jass  
Donald R. Keller  
Sergio Miranda  
Karl J. Reinhard

with support from the

Arizona State Office,  
Bureau of Land Management

BILBY RESEARCH CENTER OCCASIONAL PAPERS NO. 1

© 2002 by the Ralph M. Bilby Research Center, Northern Arizona University, Flagstaff.  
All rights reserved.

ISBN 0-9718786-3-3

## ACKNOWLEDGMENTS

The test excavation of Bighorn Cave was jointly supported by the Bureau of Land Management, Northern Arizona University, and the Museum of Northern Arizona. Individuals at these institutions who deserve credit for having helped this cooperative venture get off the ground include Don Simonis and Gary Stumpf of the BLM, Larry Agenbroad and Henry Hooper of NAU, and Don Weaver of MNA. We are especially grateful to Don Simonis, without whom the project wouldn't have happened. He was the motivating force behind the endeavor; he helped organize the field logistics, participated in much of the excavation, found money for various analyses, and generally helped out in many other ways. Henry Hooper provided research assistantships for several students working on the analysis of artifacts and faunal remains. We are indebted to the field personnel listed in Chapter 1, nearly all of whom were volunteers. We especially thank those who participated for extended intervals: most notably Peter Bungart, Mark Hovezak, Larry McConnell, and Bill Welsh. The impetus for this final report on the project came from John Rose, who followed Don Simonis as the BLM's Kingman Resource Area archaeologist. Without John's encouragement and support, the project draft manuscript might have languished for another decade and perhaps this final report would never have been completed. We appreciate the help of past and present Museum of Northern Arizona staff with curation and collections access. The report's final illustrations, excluding the projectile point illustrations, were done by Ron Redsteer; he also drew the cover art. Marvin Goad illustrated the points. Report photographs were printed by Don Keller, with the conventional images then scanned and reformatted by Zach Huntington. Louella Holter's contributions to this final report are many: technical editor, word processor, and assembler of text, tables, and graphics into a print file. Thanks, Louella, for your excellent effort. It shows in the final product.

## THE RALPH M. BILBY RESEARCH CENTER

The Ralph M. Bilby Research Center was founded in 1982 to promote interdisciplinary research at Northern Arizona University. The Bilby Research Center has been affiliated with the Center for Sustainable Environments since 1999, when the CSE was founded. The Bilby Occasional Papers Series provides an outlet for quality research on a wide range of topics. For more information please contact us at Box 6013, Northern Arizona University, Flagstaff, 86011.

## Chapter 7

### MULTIDISCIPLINARY COPROLITE ANALYSIS

*Karl J. Reinhard, Dennis R. Danielson, Mark Daniels, and Sergio Miranda*

Coprolite analysis, as reviewed by Reinhard and Bryant (1992), contributes unique and detailed information regarding diet and parasitic disease. We present here an analysis of dietary components of coprolites from Bighorn Cave using macroscopic remains, pollen concentrations, and phytoliths. In addition, we analyzed Bighorn Cave coprolites for evidence of parasitic organisms, especially intestinal worms. Such analyses of coprolites have become important methods for reconstructing past dietary and medicinal practices. Pollen concentration and phytolith quantification techniques have recently been developed, but until this report no known attempt has been made to synthesize pollen, macroscopic, and phytolith data from a single coprolite series.

#### MATERIALS AND METHODS

##### The Specimens

We analyzed 21 coprolites, designated Samples 1–21, from Components I–III of Bighorn Cave (Table 7.1). After photographing each specimen with color and black-and-white film, we made specific observations about the shape, size, and content (as evident from the surface) of each coprolite. Based on these observations we made a preliminary determination of whether the coprolites were of human origin. We also noted evidence of insect activity, such as the presence of larvae cases and beetle and arthropod holes. The preliminary examination of the coprolites indicated that all were in the range of variation for human feces. Confirmation of fecal

origin was made during the rehydration process when it was noted that bile pigments were produced by all coprolites in contact with trisodium phosphate. Sample 14, however, had an unusual green tinge to the rehydration fluid and a pronounced “musty” smell, indicating that this was not a human coprolite.

##### Pollen Analysis Methods

Two portions were broken from each coprolite: a 10 g piece for macrobotanical and parasite analyses, and a 2 g piece for pollen analysis. The 2 g pieces were fragmented and rehydrated for 24 hours in trisodium phosphate along with four *Lycopodium* tablets (each containing approximately 11,200 spores). After rehydration, each sample was treated for 30 minutes in 10% potassium hydroxide and then disaggregated with a magnetic stirrer. The potassium hydroxide softens plant tissue in the coprolite and the magnetic stirrer releases pollen that might otherwise be trapped in the plant remains. The samples were then screened through a 300 micrometer mesh. The fluid passing through the screen was collected in a large glass beaker and then centrifuged in 100 ml tubes. The concentrated solid microscopic remains were then transferred to 50 ml tubes for chemical extraction.

The extraction of the pollen involved several stages. The sediments were first treated with hydrochloric acid (70%) and were then rinsed with water. They were then bathed in nitric acid for 2 hours followed by a water

Table 7.1. Feces analyzed from Bighorn Cave (some specimen weights not recorded).

Sample No.	FS No.	Bag No.	Item No.	Weight	Comp.	Pollen	Phyto-lith	Macro	Parasite
1	177	1	1	31.3	III	x	x	x	x
2	345	2	1	27.9	III	x	x	x	x
3	345	3	1	19.7	III	x	x	x	x
4	345	4	1	17.8	III	x	x	x	x
5	345	5	1	19.4	III	x	x	x	x
6	345	6	1	21.3	III	x	x	x	x
7	345	7	1	16.2	III	x	x	x	x
8	266	1	2	48.7	II	x	x	x	x
9	390	1	1	59.2	I	x	x	x	x
10	321	1	1	—	III	x	x	x	x
11	321	1	2	—	III	x	x	x	x
12	321	1	4	—	III	—	x	x	—
13	321	1	5	—	III	x	x	x	x
14	279	6	1	—	III	x	x	x	x
15	382	1	1	16.7	II	x	x	x	x
16	382	4	1	26.0	II	x	x	x	x
17	229	1	1	—	II	x	x	x	x
18	229	1	5	—	II	x	—	x	x
19	177	2	1	12.2	III	x	—	x	x
20	177	4	1	19.4	III	x	—	x	x
21	177	3	1	24.7	III	x	—	x	x

Note: Number 14 is carnivore, not human.

rinse. Hydrofluoric acid (30%) was added and the tubes were placed in a boiling water bath for an hour, after which the samples were rinsed with water and then rinsed twice with glacial acetic acid preparatory to acetolysis. The acetolysis mixture of 1 part sulphuric acid and 9 parts acetic anhydride was added to each of the tubes, which were then placed in a boiling water bath for 40 minutes. Finally, the samples were washed with acetic acid and then several times with water (until the supernatant was clear).

Following the chemical extraction, the residue was washed with 95% alcohol and transferred to small vials in alcohol. Microscope slides were prepared by pipetting a drop of residue onto a slide, allowing most of the alcohol to evaporate, and mixing in a drop of glycerol. Then a cover glass was placed on top and sealed with fingernail polish. We examined the slides under 400x magnification, making a minimum count of 200 pollen grains for each slide. Pollen identification was aided by the comparative reference collection on file with the Palynology

Laboratory, Texas A&M University. Often-times identifications could not be made with certainty; these are followed by "cf." Pollen concentrations (Table 7.2) were determined by calculating the ratio of pollen to the known number of *Lycopodium* spores per gram.

#### Macrobotanical and Phytolith Analysis Methods

The 10 g fragments of the coprolites were rehydrated in 10% trisodium phosphate for 48 hrs as described by Fry (1977). The samples were then agitated to disaggregate them and the slurry was poured through a 500 micrometer mesh screen. The material left on top of the screen was dried on blotter paper, and the water was collected in a beaker and centrifuged to concentrate the sediments, which were saved for parasite analysis.

The dried screen remains were then examined with a binocular dissecting microscope. The macroscopic plant constituents were identified using the comparative col-

Table 7.2. Pollen concentration values for all samples, with number of taxa found.

Sample No.	Pollen Concentration	No. of Taxa
1	14,300	21
2	16,100	22
3	29,400	24
4	72,300	14
5	26,200	21
6	26,000	27
7	114,900	8
8	150,000	13
9	29,300	18
10	36,300	18
11	224,000	8
13	53,300	21
14*	20,500	19
15	4,972,800	1
16	1,136,800	14
17	17,100	15
18	129,000	19
19	>5,000,300	2
20	2,240,000	12
21	2,340,800	10

\*Non-human.

lection on file with the Ethnobotany Laboratory, Texas A&M University. In certain instances, wet mounts were made of plant tissue so the cellular and phytolith arrangements could be examined with the compound microscope.

The phytoliths were extracted following the methods of Danielson and Reinhard (1998). A small fragment from each coprolite was placed in a 500 ml beaker. We added about 5 ml of 50% hydrogen peroxide and a few crystals of potassium dichromate. The subsequent reaction destroyed all nonmineral remains and left calcium oxalate phytoliths and silica phytoliths. Identification of phytoliths was based on a reference collection of phytoliths in the University of Nebraska Microfossil Research Facility.

## RESULTS

Table 7.3 gives pollen counts, Table 7.4 presents phytolith percentages, and Table 7.5 provides macroplant remains by weight. This information is summarized below by FS number according to depositional com-

ponents. Much of the pollen reflects only the environmental pollen rain that in most specimens is dominated by Poaceae, low-spine Asteraceae, and *Quercus* (see Table 7.6 for common names of the identified taxa). Other coprolites, however, contain pollen types that certainly had a dietary origin. It is important to note that no cultivated plants are indicated by the pollen, phytolith, or macrobotanical remains. *Cucurbita* phytoliths occur in low frequency in Sample 9 but they are consistent with the wild species *C. foetidissima*.

### Component III, Formative

#### FS 177

**Pollen.** Four separate specimens were analyzed from this provenience: Samples 1 and 19–21. Three of these have an abundance of dietary pollen but the fourth contains what is probably environmental pollen. Mustard pollen (Brassicaceae) dominates Sample 19 to the near exclusion of all other types. *Ephedra* pollen dominates Samples 20 and 21.

In areas where *Ephedra* is common, its pollen in soils varies between 8 and 45 percent (Reinhard, unpublished counts of modern ecosystems). The variation in count depends on whether the sample was collected in an area devoid of small forbs where high *Ephedra* counts occur or in areas where forbs are abundant. The amount of *Ephedra* pollen in the coprolites clearly exceeds that expected in natural pollen rain. Pollen concentration results (Table 7.2) show that Samples 19, 20, and 21 contain quantities in the millions of pollen grains per gram. Sample 19 contains such an abundance of pollen that definitive quantification was impossible other than that pollen is in excess of 5,000,000 grains per gram. Such large quantities of pollen could have been ingested with tea brewed from *Ephedra*.

Sample 1 contains about 14,300 pollen grains per gram. These grains are in a poor state of preservation. The count reflects a natural spectrum of predominantly wind-borne pollen types such as *Quercus*, low-spine Asteraceae, Poaceae, and Cheno-Am.



Table 7.3. Pollen counts for fecal samples from Bighorn Cave; number in parentheses is column percentage rounded to the nearest tenth. (Sample 12 contained no pollen.)

Taxa	1	2	3	4	5
<i>Acacia</i>	6 (2.5)	1 (0.4)	—	—	—
<i>Acer</i>	—	—	1 (0.5)	—	—
<i>Acer negundo</i> cf.	—	—	—	—	—
<i>Alnus</i>	—	—	—	—	—
<i>Artemisia</i>	1 (0.4)	1 (0.4)	5 (2.3)	1 (0.5)	3 (1.5)
Low spine	57 (24.1)	49 (21.9)	28 (12.7)	38 (19.0)	42 (20.5)
High spine	8 (3.4)	4 (1.8)	5 (2.3)	6 (3.0)	6 (2.9)
<i>Astragalus</i>	—	—	—	—	—
Berberidaceae	—	1 (0.4)	1 (0.5)	6 (3.0)	—
Brassicaceae	—	—	1 (0.5)	—	2 (1.0)
<i>Carex</i>	9 (3.8)	6 (2.7)	—	—	—
<i>Cercidium</i>	—	—	1 (0.5)	—	—
Cheno-Am	27 (11.4)	6 (2.7)	5 (2.3)	6 (3.0)	4 (2.0)
<i>Ephedra</i> sp.	—	—	—	11 (5.5)	—
<i>Ephedra viridis</i> cf.	7 (3)	—	—	—	5 (2.4)
<i>Ephedra nevadensis</i> cf.	—	—	—	—	—
Ericaceae cf.	—	—	1 (0.5)	—	—
<i>Eriogonum</i> cf.	5 (2.1)	3 (1.3)	3 (1.4)	—	—
Euphorbiaceae	—	—	—	—	—
Fabaceae	1 (0.4)	—	3 (1.4)	—	1 (0.5)
<i>Fraxinus</i>	—	—	—	—	—
<i>Juniperus</i>	—	7 (3.1)	2 (0.9)	—	2 (1.0)
Labiatae	—	2 (0.9)	—	—	—
<i>Larrea</i>	3 (1.3)	—	4 (1.8)	—	2 (1.0)
Ligulifloreae	—	—	—	—	1 (0.5)
Liliaceae	—	—	1 (0.5)	—	—
<i>Opuntia</i>	—	—	—	—	—
Onagraceae	—	—	—	—	—
<i>Physalis</i>	1 (0.4)	—	—	—	—
<i>Pinus</i>	10 (4.2)	12 (5.4)	3 (1.4)	13 (6.5)	1 (0.5)
<i>Ptelea</i> cf.	—	—	—	—	—
Poaceae	28 (11.8)	59 (26.5)	79 (35.7)	28 (14)	51 (24.9)
<i>Polygonum</i> cf.	1 (0.4)	—	—	—	—
<i>Populus</i>	1 (0.4)	1 (0.4)	10 (4.5)	—	—
<i>Portulaca</i>	—	—	—	—	—
<i>Prosopis</i>	2 (0.8)	—	—	—	1 (0.5)
<i>Prunus</i> cf.	—	—	1 (0.5)	—	—
<i>Ptelia</i>	—	—	—	—	—
<i>Quercus</i>	21 (8.9)	21 (9.4)	31 (14.0)	29 (14.5)	22 (10.7)
Rhamnaceae	—	2 (0.9)	2 (0.9)	—	2 (1.0)
Rosaceae	—	—	—	2 (1.0)	—
Rubiaceae	—	1 (0.4)	—	—	—
Salicaceae	—	—	—	—	—
<i>Salix</i>	9 (3.8)	4 (1.8)	8 (3.6)	3 (1.5)	3 (1.5)
probable <i>Salix</i>	—	—	—	—	—
Solanaceae	—	—	—	—	—
<i>Solanum</i> cf.	—	—	—	—	—
<i>Sphaeralcea</i>	—	—	—	—	—
<i>Trifolium</i> cf.	—	—	—	—	—
<i>Typha latifolia</i>	2 (0.8)	1 (0.4)	—	—	6 (2.9)
Unbelliferae	—	—	1 (0.5)	—	—
Unidentifiable	37 (15.6)	37 (16.6)	17 (7.7)	8 (4.0)	8 (3.9)
Unknown	—	—	—	1 (0.5)	1 (0.5)
Verbenaceae	—	1 (0.4)	8 (3.6)	—	1 (0.5)
<i>Yucca</i>	1 (0.4)	—	—	48 (24)	41 (20)

Table 7.3 (continued)

Taxa	6	7	8	9	10
<i>Acacia</i>	—	—	—	—	—
<i>Acer</i>	—	—	—	—	—
<i>Acer negundo</i> cf.	—	—	—	—	—
<i>Alnus</i>	1 (0.4)	—	—	—	—
<i>Artemisia</i>	4 (1.6)	—	—	1 (0.5)	6 (2.8)
Low spine	66 (27.0)	6 (3.0)	20 (10.0)	54 (27.0)	61 (28.8)
High spine	23 (9.4)	2 (1.0)	19 (9.5)	7 (3.5)	6 (2.8)
<i>Astragalus</i>	—	—	—	1 (0.5)	—
Berberidaceae	—	—	—	—	—
Brassicaceae	1 (0.4)	5 (2.5)	5 (2.5)	7 (3.5)	3 (1.4)
<i>Carex</i>	1 (0.4)	—	—	—	—
<i>Cercidium</i>	—	—	—	—	—
Cheno-Am	1 (0.4)	160 (80.0)	4 (2.0)	9 (4.5)	6 (2.8)
<i>Ephedra</i> sp.	—	—	—	1 (0.5)	—
<i>Ephedra viridis</i> cf.	—	—	—	—	—
<i>Ephedra nevadensis</i> cf.	—	—	—	4 (2.0)	4 (2.0)
Ericaceae cf.	—	—	—	—	—
<i>Eriogonum</i> cf.	10 (4.1)	—	—	1 (0.5)	—
Euphorbiaceae	1 (0.4)	—	—	—	1 (0.5)
Fabaceae	1 (0.4)	—	—	—	—
<i>Fraxinus</i>	1 (0.4)	—	—	—	—
<i>Juniperus</i>	1 (0.4)	—	—	1 (0.5)	—
Labiatae	—	—	—	—	—
<i>Larrea</i>	7 (2.9)	—	—	—	8 (3.8)
Ligulafloreae	—	—	—	—	—
Liliaceae	—	—	—	—	—
<i>Opuntia</i>	—	—	—	—	—
Onagraceae	—	—	—	—	—
<i>Physalis</i>	—	—	—	—	—
<i>Pinus</i>	8 (3.3)	—	1 (0.5)	11 (5.5)	10 (4.7)
<i>Ptelea</i> cf.	1 (0.4)	—	—	—	—
Poaceae	27 (11.1)	—	8 (4.0)	7 (3.5)	18 (8.5)
<i>Polygonum</i> cf.	—	—	—	—	—
<i>Populus</i>	7 (2.9)	8 (4.0)	1 (0.5)	—	9 (4.2)
<i>Portulaca</i>	1 (0.4)	—	—	—	—
<i>Prosopis</i>	—	—	—	—	1 (0.5)
<i>Prunus</i> cf.	—	—	—	—	—
<i>Ptelia</i>	—	—	—	—	—
<i>Quercus</i>	47 (19.3)	2 (1.0)	100 (50.0)	30 (15.0)	52 (24.5)
Rhamnaceae	4 (1.6)	—	—	—	—
Rosaceae	—	—	—	—	—
Rubiaceae	—	—	—	—	—
Salicaceae	—	—	—	—	—
<i>Salix</i>	5 (2.0)	—	1 (0.5)	29 (14.5)	3 (1.4)
probable <i>Salix</i>	—	—	—	21 (10.5)	—
Solanaceae	—	—	—	—	—
<i>Solanum</i> cf.	—	—	—	—	—
<i>Sphaeralcea</i>	1 (0.4)	—	—	—	—
<i>Trifolium</i> cf.	—	—	—	1 (0.5)	—
<i>Typha latifolia</i>	1 (0.4)	—	2 (1.0)	—	—
Unbelliferae	—	—	1 (0.5)	—	—
Unidentifiable	19 (7.9)	16 (8.0)	34 (17.0)	14 (7.0)	20 (9.4)
Unknown	—	—	4 (2.0)	1 (0.5)	1 (0.5)
Verbenaceae	1 (0.4)	1 (0.5)	—	—	2 (0.9)
<i>Yucca</i>	3 (1.2)	—	—	—	1 (0.5)

Table 7.3 (continued)

Taxa	11	13	14	15	16
<i>Acacia</i>	—	—	1 (0.5)	—	—
<i>Acer</i>	—	—	1 (0.5)	—	—
<i>Acer negundo</i> cf.	1 (0.5)	—	—	—	—
<i>Alnus</i>	—	—	—	—	—
<i>Artemisia</i>	—	—	1 (0.5)	—	1 (0.5)
Low spine	3 (1.5)	38 (19.0)	28 (14.0)	—	28 (14.0)
High spine	—	2 (1.0)	9 (4.5)	—	6 (3.0)
<i>Astragalus</i>	—	—	—	—	—
Berberidaceae	—	—	—	—	—
Brassicaceae	16 (8.0)	9 (4.5)	—	—	2 (1.0)
<i>Carex</i>	—	—	—	—	—
<i>Cercidium</i>	—	1 (0.5)	—	—	—
Cheno-Am	—	1 (0.5)	29 (14.5)	—	13 (6.5)
<i>Ephedra</i> sp.	—	2 (1)	3 (1.5)	—	1 (0.5)
<i>Ephedra viridis</i> cf.	—	—	—	—	—
<i>Ephedra nevadensis</i> cf.	—	—	—	—	—
Ericaceae cf.	—	—	—	—	—
<i>Eriogonum</i> cf.	—	3 (1.5)	1 (0.5)	—	—
Euphorbiaceae	—	—	—	—	—
Fabaceae	1 (0.5)	1 (0.5)	1 (0.5)	—	—
<i>Fraxinus</i>	—	—	—	—	—
<i>Juniperus</i>	2 (1.0)	—	6 (3.0)	—	—
Labiatae	—	1 (0.5)	—	—	—
<i>Larrea</i>	—	9 (4.5)	8 (4.0)	—	—
Ligulifloreae	—	—	1 (0.5)	—	—
Liliaceae	—	—	1 (0.5)	—	6 (3.0)
<i>Opuntia</i>	—	53 (26.5)	—	—	—
Onagraceae	—	—	—	—	1 (0.5)
<i>Physalis</i>	—	—	—	—	—
<i>Pinus</i>	2 (1.0)	—	26 (13.0)	—	3 (1.5)
<i>Ptelea</i> cf.	—	—	—	—	—
Poaceae	—	36 (18.0)	39 (18.5)	—	7 (3.5)
<i>Polygonum</i> cf.	—	—	—	—	—
<i>Populus</i>	—	—	—	—	1 (0.5)
<i>Portulaca</i>	—	—	—	—	—
<i>Prosopis</i>	—	—	—	—	—
<i>Prunus</i> cf.	—	—	—	—	—
<i>Ptelia</i>	—	—	—	—	—
<i>Quercus</i>	3 (1.5)	16 (8.0)	30 (15.0)	—	—
Rhamnaceae	—	3 (1.5)	—	—	—
Rosaceae	—	—	—	—	—
Rubiaceae	—	—	—	—	—
Salicaceae	—	—	—	—	—
<i>Salix</i>	172 (86.0)	2 (1.0)	1 (0.5)	222 (100.0)	121 (60.5)
probable <i>Salix</i>	—	—	—	—	—
Solanaceae	—	1 (0.5)	1 (0.5)	—	—
<i>Solanum</i> cf.	—	2 (1.0)	—	—	—
<i>Sphaeralcea</i>	—	—	—	—	—
<i>Trifolium</i> cf.	—	—	—	—	—
<i>Typha latifolia</i>	—	1 (0.5)	—	—	2 (1.0)
Unbelliferae	—	—	—	—	—
Unidentifiable	—	15 (7.5)	12 (6.0)	—	8 (4.0)
Unknown	—	1 (0.5)	—	—	—
Verbenaceae	—	—	—	—	—
<i>Yucca</i>	—	3 (1.5)	—	—	—

Table 7.3 (continued)

Taxa	17	18	19	20	21
<i>Acacia</i>	—	—	—	—	—
<i>Acer</i>	1 (0.5)	—	—	—	—
<i>Acer negundo</i> cf.	—	—	—	—	—
<i>Alnus</i>	—	—	—	—	—
<i>Artemisia</i>	1 (0.5)	1 (0.5)	—	—	—
Low spine	21 (9.7)	35 (16.4)	—	4 (2.0)	12 (5.7)
High spine	5 (2.3)	4 (1.9)	—	1 (0.5)	1 (0.5)
<i>Astragalus</i>	—	—	—	—	—
Berberidaceae	—	—	—	—	—
Brassicaceae	1 (0.5)	9 (14.2)	199 (99.5)	6 (3.0)	4 (1.9)
<i>Carex</i>	—	—	—	—	—
<i>Cercidium</i>	—	—	—	—	—
Cheno-Am	42 (19.4)	9 (4.2)	—	1 (0.5)	3 (1.4)
<i>Ephedra</i> sp.	3 (1.4)	4 (1.9)	—	149 (74.5)	170 (81.3)
<i>Ephedra viridis</i> cf.	—	4 (1.9)	—	—	—
<i>Ephedra nevadensis</i> cf.	—	—	—	—	—
Ericaceae cf.	—	—	—	—	—
<i>Eriogonum</i> cf.	—	1 (0.5)	—	—	—
Euphorbiaceae	—	—	—	—	—
Fabaceae	1 (0.5)	1 (0.5)	—	—	—
<i>Fraxinus</i>	—	—	—	—	—
<i>Juniperus</i>	—	—	—	—	—
Labiatae	—	—	—	—	—
<i>Larrea</i>	1 (0.5)	—	—	5 (2.5)	4 (1.9)
Ligulafloreae	—	—	—	—	2 (1.0)
Liliaceae	—	—	—	—	—
<i>Opuntia</i>	5 (2.3)	—	—	—	—
Onagraceae	—	1 (0.5)	—	—	—
<i>Physalis</i>	—	—	—	—	—
<i>Pinus</i>	3 (1.4)	12 (5.6)	—	1 (0.5)	—
<i>Ptelea</i> cf.	—	—	—	—	—
Poaceae	88 (40.7)	102 (47.9)	1 (0.5)	18 (9.0)	5 (2.4)
<i>Polygonum</i> cf.	—	—	—	—	—
<i>Populus</i>	—	4 (1.9)	—	—	—
<i>Portulaca</i>	—	—	—	—	—
<i>Prosopis</i>	—	1 (0.5)	—	—	—
<i>Prunus</i> cf.	—	—	—	—	—
<i>Ptelia</i>	—	—	—	—	—
<i>Quercus</i>	15 (6.9)	14 (6.6)	—	1 (0.5)	1 (0.5)
Rhamnaceae	—	1 (0.5)	—	—	—
Rosaceae	—	—	—	—	—
Rubiaceae	—	—	—	—	—
Salicaceae	—	—	—	—	—
<i>Salix</i>	22 (10.2)	4 (1.9)	—	1 (0.5)	—
probable <i>Salix</i>	—	—	—	—	—
Solanaceae	—	—	—	—	—
<i>Solanum</i> cf.	—	—	—	—	—
<i>Sphaeralcea</i>	—	—	—	—	—
<i>Trifolium</i> cf.	—	1 (0.5)	—	—	—
<i>Typha latifolia</i>	—	—	—	—	—
Unbelliferae	—	—	—	—	—
Unidentifiable	7 (3.2)	5 (2.3)	—	12 (6.0)	7 (3.3)
Unknown	—	—	—	—	—
Verbenaceae	—	—	—	—	—
<i>Yucca</i>	—	—	—	1 (0.5)	—

Table 7.4. Phytolith percentages for fecal samples from Bighorn Cave. The numbers are percentage expressions of the total count for each coprolite. The total phytoliths counted are also presented. The *Cucurbita* phytoliths are consistent with the species *Cucurbita foetidissima*.

Taxa	1	2	3	4	5	6	7	8	9	10
Agavaceae	55	68.5	77.5	1	–	25	0.9	43	15	50
Cactaceae	14	0.3	4	50	60	26.5	1	–	15	–
Cheno-Am	6	0.7	–	4	37	34	5	–	6	–
Chloridoid	–	–	0.7	–	–	–	–	2	0.4	–
Festucoid	–	5	2	2	–	2	–	10	5	50
Panicoid	–	0.3	0.7	0.9	–	0.8	–	0.4	0.8	–
Other grass	–	–	–	–	–	–	–	0.7	–	–
Fabaceae	25	25	15	42	3	12	93	43	69	–
<i>Cucurbita</i>	–	–	–	–	–	–	–	–	9	–
Total count	275	288	281	224	229	240	226	268	225	10

Table 7.4, continued

Taxa	11	12	13	14	15	16	17
Agavaceae	98	88	6	6	2.8	22	34
Cactaceae	2	–	28	30	43	51	44
Cheno-Am	–	0.5	29	9	34	23	20
Chloridoid	–	1	–	–	–	–	–
Festucoid	–	4	0.4	2	3	3	0.4
Panicoid	–	6	–	0.7	12	0.4	1
Other grass	–	–	–	–	–	–	–
Fabaceae	–	–	36	52	5	–	–
<i>Cucurbita</i>	–	–	–	–	–	–	–
Total count	49	209	238	286	264	225	280

The poor preservation suggests that the pollen types were in the environment for a long time prior to human consumption; these grains may have been consumed with drinking water.

**Macro and Phytolith.** Sample 1 contains small amounts of seed. One seed appears to be of the genus *Asclepias*. The others are mustard seeds similar in morphology to *Cenchrus*. Charcoal is present in very small amounts. The vast majority of the macrofossil remains consist of short screwbean mesquite fibers arranged in tiny concentric circles. Most often the circular fibers occur separately, but in some instances the fibers are held in place with a thin, rough cuticle. Spiral-shaped arrangements of the fiber, resembling tendrils, are also present. These fibers originated from the consumption of

screwbean mesquite, thickets of which grow along the washes of the Black Mountains. The phytoliths show that other plants were also eaten. Agavaceae accounts for 55 percent of the phytoliths and Cactaceae and Fabaceae account for 14 and 25 percent respectively. Therefore, three sources of fiber are indicated by the phytoliths.

Sample 19 contains small bone fragments. As in most other Bighorn Cave human coprolites, the bone fragments are very small and include fragments of trabecular bone, which suggests that the bone was ground for consumption. The majority of the macroscopic remains from this coprolite consist of *Opuntia* epidermis and fiber. Druse crystals are also present, sometimes still in place inside of epidermal fragments. Also present is a *Quercus* leaf and a thorn.

Table 7.5. Macroscopic remains for the Bighorn Cave fecal specimens by weight in grams. Components weighing less than 0.01 grams are indicated by "t".

[illegible]

Table 7.5 continued

[illegible]

Table 7.6. Scientific and common names of pollen taxa recovered from coprolites at Bighorn Cave.

Scientific Name	Common Name
<i>Acacia</i>	Acacia
<i>Acer</i>	Maple
<i>Alnus</i>	Alder
<i>Artemisia</i>	Sage
Asteraceae	Composite family
<i>Astragalus</i>	Vetch
Berberidaceae	Barberry family
Brassicaceae	Mustard family
<i>Carex</i>	Sedge
<i>Cercidium</i>	Palo verde
Cheno-Am	Goosefoot family and amaranth (pigweed) family
<i>Ephedra</i> sp.	Mormon tea, species unknown
Ericaceae	Heather family
<i>Eriogonum</i>	Wild buckwheat
Euphorbiaceae	Spurge family
Fabaceae	Legume family
<i>Fraxinus</i>	Ash
High spine	Asteraceae subgroup that includes sunflower
<i>Juniperus</i>	Juniper
Labiatae	Mint family
<i>Larrea</i>	Creosote
Ligulifloreae	Asteraceae subgroup that includes dandelion
Liliaceae	Lily family
Low spine	Asteraceae subgroup that includes ragweed
<i>Opuntia</i>	Prickly pear
Onagraceae	Evening primrose family
<i>Physalis</i>	Ground cherry
<i>Pinus</i>	Pine
Poaceae	Grass family
<i>Polygonum</i>	Smartweed
<i>Populus</i>	Cottonwood
<i>Portulaca</i>	Portulaca
<i>Prosopis</i>	Mesquite
<i>Ptelia</i>	Hop tree
<i>Quercus</i>	Oak
Rhamnaceae	Buckthorn family
Rubiaceae	Madder family
<i>Salix</i>	Willow
Solanaceae	Potato family
<i>Sphaeralcea</i>	Mallow
<i>Trifolium</i> type	Clover-like legume
<i>Typha latifolia</i>	Cattail
Unbelliferae	Carrot family
Unidentifiable	Pollen grains too degraded to identify certainly
Unknown	Pollen grain morphology could not be identified
Verbenaceae	Vervain family
<i>Yucca</i>	Yucca

Sample 20 also contains small bone fragments, apparently from fish. Also present are remains of *Allium* (onion) bulbs. The majority of the coprolite consists of a fine brown substance of unknown origin. A wood fragment is also present. Sample 21 contains bone similar to Sample 19. Some fiber is present, as is the cuticle of an unidentified plant.

#### FS 279

The single coprolite from this provenience is from a nonhuman carnivore. Before rehydration, it was composed of a hard, dark crust surrounding a core of bone and air. It contains about 20,500 pollen grains per gram. The dominant types are windborne. The pollen content of this coprolite no doubt mirrors the environmental pollen rain present at the time of defecation.

The macroscopic part of Sample 14 consists entirely of bone and hair. The bones are those of a small rodent. The plant dietary component of this coprolite is represented in the phytolith count: Fabaceae is most common, probably from mesquite, followed by Cactaceae, Cheno-Am, Agavaceae, and various grasses.

#### FS 321

**Pollen.** Three coprolites (10, 11, and 13) were processed from this provenience. An additional coprolite (12) was analyzed, but processing and reprocessing recovered no pollen. The pollen analysis from 10 did not demonstrate any clear examples of dietary types. However, Sample 11 is dominated by willow (*Salix*) and 13 contains a relatively large percentage of *Opuntia* pollen. *Salix* is a windborne type that is especially common in spring. However, it is improbable that *Salix* would normally make up 86 percent of the pollen rain. This undoubtedly reflects dietary use of the plant. *Opuntia* rarely accounts for more than 2 percent of the normal pollen rain, even in a prickly pear patch (Reinhard, unpublished pollen counts of modern ecological zones). The more than 25 percent *Opuntia* pollen in Sample 13 is clearly

a result of dietary usage of the plant. In Sample 10, 36,300 pollen grains per gram are present, and 53,300 grains per gram are present in 13. A relatively high pollen concentration value of 224,000 grains per gram was obtained from Sample 11.

**Macro and Phytolith.** Sample 10 consists of small bone fragments similar to those found in 19. Also present is a small amount of *Opuntia* epidermis. Most of the coprolite consists of fiber. Very few phytoliths were found; half are from a cactus and half are from a festucoid grass. Samples 11 and 12 consist solely of fiber, perhaps *Agave*. Samples 11 and 12 both have abundant phytoliths from an Agavaceae plant, consistent with *Agave*. Sample 13 yielded mostly mustard seed. The seeds are very abundant and seem to have been eaten as a "cake." Many seeds are broken and may have been prepared by crushing or grinding. The seed morphology is very close to that of *Descurainia*. Also present in this sample are the circular fibers from screwbean mesquite described for Sample 1. Cactaceae, Cheno-Am, and Fabaceae phytoliths are very common in this sample, along with small amounts of Agavaceae.

#### FS 345

**Pollen.** Six coprolites were analyzed from FS 345 (Samples 2–7). Two of these (2 and 6) were different from the other four in having poorly preserved pollen similar to Sample 1 of FS 177. Both of these samples probably reflect the natural pollen rain. The relatively high Poaceae count of Sample 3 suggests that Poaceae pollen was introduced into the intestine through dietary means, but this cannot be said with certainty. *Yucca* pollen in 5 and 6 clearly indicates a dietary use of this plant because *Yucca* pollen is not normally present in pollen rain in such high percentages. The dominance of Cheno-Am pollen in Sample 7 probably has a dietary origin; although Cheno-Am is a windblown type, its percentage in Sample 7 exceeds the natural pollen rain of Cheno-Am that might be incidentally ingested.



**Macro and Phytolith.** Sample 2 contains an abundance of screwbean mesquite circular fibers. Also present are *Chenopodium* seeds and possibly *Panicum* seed. These seeds appear unground and were probably eaten whole. Sample 3 contains what appear to be well-masticated insect fragments. Fiber dominates the macrofossil remains, but *Sporobolus* seeds are also present in small numbers. Samples 2 and 3 are both dominated by Agavaceae phytoliths with a strong showing of Fabaceae phytoliths. Sample 4 consists mostly of coarse fiber. *Allium* tissue is also present, as is a possible *Yucca* seed. The phytolith analysis shows that large amounts of cactus and Fabaceae were eaten. Sample 5 consists mostly of *Opuntia* epidermis and some fiber. The phytoliths from Sample 5 are dominated by Cactaceae, with some Cheno-Am. Sample 6 consists mostly of *Opuntia* epidermis and some fiber. Sample 7 consists primarily of circular screwbean mesquite fibers, along with a few *Sporobolus* seeds that appear to be parched. Nearly all the phytoliths are from Fabaceae.

#### Component II, Terminal Archaic

##### FS 229

**Pollen.** Both of the samples from FS 229 have high percentages of Poaceae pollen. Although this is a windborne type and consequently is easily ingested by inhalation and drinking, as well as eating, the high percentages of Samples 17 and 18 indicate that it was a dietary type in these incidences. The average Poaceae percentage for all human coprolites from Bighorn Cave is 13.8 percent. Sample 17 contains 41 percent and 18 contains 48 percent. This strongly suggests that grass in these instances is dietary. Another dietary type, Brassicaceae, is represented by a 14 percent rate in Sample 18. This is a high percentage for an insect-pollinated species, and probably indicates a dietary origin. Both 17 and 18 contain large amounts of fine charcoal in the pollen preparations. This is similar to coprolites from Dust Devil Cave that we have examined in which high grass pollen percentages occurred with large amounts of charcoal. We

suspect that parching grass seeds or inflorescences accounts for the co-occurrence. Pollen concentration indicates that 17,000 and 129,000 grains per gram are present in 17 and 18, respectively (Table 7.2).

**Macro and Phytolith.** In Samples 17 and 18, nearly all of the rehydrated material passed through the screen. A juniper nut and a bit of fine fiber are present in Sample 17. Only sand and gravel were present after 18 was processed. The gravel and sand were probably incorporated in the coprolite from contact with the cave substrate. Considering the pollen analyses, it is likely that ground grass seed, perhaps parched, constituted the meals represented in Samples 17 and 18. However, the phytoliths show that at least three other plants were eaten: Agavaceae, Cactaceae, and Cheno-Am.

##### FS 266

**Pollen.** The one coprolite from this provenience contained about 150,000 pollen grains per gram. The pollen counts are dominated by *Quercus* (50%). The relatively high percentage of unidentifiable pollen (17%) constitutes spindled or crushed grains that are probably *Quercus*. In scrubland dominated by *Quercus*, up to 45 percent of the natural pollen rain is derived from it (Reinhard, unpublished pollen counts of modern ecological zones). The average *Quercus* percentage for all human coprolites from Bighorn Cave is 10.1 percent. The high percentage in Sample 8 could be due to dietary reasons. However, *Quercus* is a prolific pollen producer, and is wind pollinated. Therefore, it is possible that the high percentage is due to incidental consumption with food or drink.

**Macro and Phytolith.** Sample 8 consists of a finely ground, dark brown meal. It is not possible to determine the source of the meal from macroscopic or palynological examination. However, the phytoliths show that Agavaceae composed a large part of the meal. Also, various grasses, including Chloroid, Festucoid, and Panicoid were eaten. Also, Fabaceae phytoliths are abundant, probably from mesquite.

## FS 382

**Pollen.** Samples 15 and 16 have very high pollen grain per gram values (4,972,800 and 1,136,000 respectively). A large amount of fine charcoal is present in 16. Both samples contain large percentages of *Salix* pollen that certainly indicate consumption of this plant. In 15, the large number of *Salix* pollen grains exceeds all other types.

**Macro and Phytolith.** Sample 15 is dominated by screwbean mesquite fibers. Minor components are *Descurainia* cf. seeds and what looks like a fragment of snakeskin. Cactus, Cheno-Am, and grass dominate the phytolith count from this sample. Sample 16 consists of smashed acorn hulls in a fine brown matrix. Agavaceae, Cactaceae, and Cheno-Am phytoliths are abundant in this sample.

## Component I, Late Archaic

## FS 390

The single sample from this provenience contains badly degraded pollen that might reflect consumption of *Salix*. Of the pollen grains counted, almost 15 percent are *Salix*. Degraded grains that are probably *Salix* constitute another 10 percent. Pollen concentration indicates 29,300 grains per gram.

The macro remains of this specimen consist solely of coarse fiber. Agavaceae, Cactaceae, and Fabaceae phytoliths are abundant.

## PARASITOLOGICAL ANALYSIS

The centrifuged sediments left over from macrobotanical separation were transferred into vials and sedimented gravitationally in acetic formalin alcohol. Previous work with this technique has demonstrated that parasite eggs and larvae settle out in the upper level of the sediment (Reinhard 1985a). Microscopic preparation requires only a few drops of sediment. Drops of sediment from the upper layers of the debris are placed on a microscope slide. When the acetic formalin alcohol has nearly evaporated, a small drop of glycerol is added to the preparation. The sediment and glycerol are thoroughly mixed with an applicator stick and a cover glass is placed on top. The cover glass is sealed with

finger nail polish and the preparation is scanned under 200x for helminth eggs. For quantification, one *Lycopodium* tablet is added to each gram of coprolite. This allows for egg per gram (epg) calculation of parasite infections by calculating the known ratio of spores to eggs or larvae.

No definite parasite infections are apparent in the 1 carnivore and 20 human coprolites. However, what are possibly tapeworm eggs of the family Taeniidae, or fungal spores resembling taeniid eggs, were found in Samples 2 (640 epg) and 10 (200 epg). The objects are spherical, averaging 32 micrometers in diameter, and exhibit a thick, radially striated wall. These observations are consistent with the form of taeniid eggs. However, no hooklets were visible on the interiors of the possible eggs. Because the hooklets are not present, it is possible that the objects are fungal and not parasitic.

Free-living, nonparasitic nematode larvae are present in Samples 9 and 16. These are the only coprophagous organisms found in the analysis. Free-living mites and other arthropods that are often found in the microscopic fraction of coprolites from the Southwest are absent in the Bighorn Cave coprolites.

## DISCUSSION

The data resulting from the analyses of three different fecal constituents are combined in Tables 7.7 and 7.8. The pollen analysis demonstrates that the prehistoric inhabitants of the cave lived in a xeric environment but utilized nearby water sources. The presence of *Acer*, *Salix*, *Populus*, *Typha*, and *Carex* pollen demonstrates the use of wet environments. The analysis shows that at least one of these genera, *Salix*, was of dietary use. An overall dry environment is indicated by *Ephedra*, *Yucca*, *Artemisia*, *Prosopis*, *Acacia*, *Larrea*, *Juniperus*, *Sphaeralcea*, and *Opuntia*. Of these plants, *Yucca*, *Opuntia*, and *Ephedra* were eaten. *Prosopis* was also consumed but the evidence for this comes from phytoliths and macroplant remains. Other dietary plants that could be from either environment include mustard, grass, and Cheno-Am.

Table 7.7. Summary of dietary remains per coprolite.

- 
- Sample 1: Phytolith data indicate consumption of plants in the Agavaceae, Cactaceae, and Fabaceae. The most likely genera consumed were *Agave*, *Opuntia*, and *Prosopis*. Phytoliths consistent with *Chenopodium* or *Amaranthus* were also found. The main macrofloral dietary component was mesquite. A small animal was eaten.
- Sample 2: Phytolith data indicate consumption of plants in the Agavaceae and Fabaceae. The most likely genera consumed were *Agave* and *Prosopis*. Grass phytoliths were also present. The main macrofloral dietary component was mesquite. Small amounts of goosefoot seed were also eaten.
- Sample 3: Phytolith data indicate consumption of plants in the Agavaceae and Fabaceae. The most likely genera consumed were *Agave* and *Prosopis*. Grass phytoliths and some cactus phytoliths were also present. Mesquite pods, prickly pear pads, and course fiber were the main macrofloral dietary components of this sample. A small animal was eaten.
- Sample 4: Pollen data indicate that yucca buds or flowers were eaten. Phytolith data indicate consumption of plants in the Cactaceae and Fabaceae. The most likely genera consumed were *Opuntia* and *Prosopis*. Small amounts of Agavaceae and grass phytoliths are also present. Mesquite pods and course fiber were the main macrofloral dietary components of this sample.
- Sample 5: Pollen data indicate that yucca buds or flowers were eaten. Phytolith data indicate that Cactaceae and Cheno-Am were eaten. The most likely genera eaten were *Opuntia* and *Chenopodium-Amaranthus*. Prickly pear pads were the main macrofloral component.
- Sample 6: Phytolith data indicate consumption of plants in the Agavaceae, Cactaceae, Cheno-Am, and Fabaceae. The most likely genera consumed were *Agave*, *Opuntia*, *Prosopis*, and *Chenopodium-Amaranthus*.
- Sample 7: Pollen data indicate that buds, flowers, or greens of species of pigweed or goosefoot were eaten. Phytoliths show that Fabaceae was consumed, most likely *Prosopis*. Small numbers of Agavaceae, Cactaceae, and Cheno-Am phytoliths were also present. The main macrofloral dietary component was mesquite.
- Sample 8: Phytolith data indicate consumption of plants in the Agavaceae and Fabaceae. The most likely genera consumed were *Agave* and *Prosopis*. Phytoliths indicate that grass was also eaten. There was no identifiable macroscopic component.
- Sample 9: Pollen data indicate that willow was eaten. Phytolith data indicate that Fabaceae, most probably *Prosopis*, was eaten. Smaller amounts of Agavaceae, Cactaceae, Cheno-Am, and grass were consumed as indicated by phytoliths. This was the one specimen with *Cucurbita* phytoliths, which were consistent with the wild gourd *C. foetidissima*. Fiber and unidentifiable plant residue make up the macrofloral dietary component. A small animal was eaten.
- Sample 10: Small numbers of phytoliths were present from Agavaceae and grass. Course fiber and prickly pear pads make up the macrofloral dietary component. A small animal was eaten.
- Sample 11: Pollen data indicate that willow was eaten. Phytoliths show that Agavaceae, probably *Agave*, was eaten. Some cactus was also eaten as indicated by phytoliths. Course fiber and some prickly pear pad make up the macrofloral dietary component.
- Sample 12: Phytoliths show that Agavaceae, probably *Agave*, was eaten. Some grass and Cheno-Am was also eaten as shown by phytoliths. Course fiber and some prickly pear pad make up the macrofloral dietary component. A small animal was eaten.
- Sample 13: Pollen data indicate that prickly pear buds or flowers were eaten. Phytolith data indicate consumption of plants in the Cactaceae, Cheno-Am, and Fabaceae families. The most likely genera consumed were *Opuntia*, *Prosopis*, and *Chenopodium-Amaranthus*. Lesser amounts of Agavaceae were consumed, as shown by phytoliths. Mesquite pods are the main macrofloral dietary component. A small animal was eaten.
- Sample 15: Pollen data indicate that willow was eaten. Phytolith data indicate consumption of plants in the Cactaceae and Cheno-Am families. The most likely genera consumed were *Opuntia* and *Chenopodium-Amaranthus*. Lesser amounts of Agavaceae and Fabaceae were consumed, as shown by phytoliths. Mesquite pods are the main macrofloral dietary component.
- Sample 16: Pollen data indicate that willow was eaten. Phytolith data indicate consumption of plants in the Agavaceae, Cactaceae, and Cheno-Am families. Genera most likely consumed were *Agave*, *Opuntia*, and *Chenopodium-Amaranthus*. The macrofloral dietary component comprises wild squash seeds.

Table 7.7 (continued)

Sample 17: Pollen data suggest that grass was eaten. Phytolith data indicate consumption of plants in the Agavaceae, Cactaceae, and Cheno-Am families. Genera most likely consumed were *Agave*, *Opuntia*, and *Chenopodium-Amaranthus*. Mesquite dominates the macrofloral remains although there are traces of prickly pear pad and traces of leaf, stem, and coarse fiber. A small animal was eaten.

Sample 18: Pollen data strongly suggest that grass was eaten. No phytolith analysis for this sample. No macrofloral remains were recovered.

Sample 19: Pollen data indicate that buds, flowers, or greens of species of mustard were eaten. No phytolith analysis for this sample. Mesquite pods dominate the macrofloral remains.

Sample 20: Pollen data indicate that Mormon tea was consumed. No phytolith analysis for this sample. The macrofloral component is composed of unidentifiable plant residue with some mesquite pod. A small animal was eaten.

Sample 21: Pollen data indicate that Mormon tea was consumed. No phytolith analysis for this sample. The macrofloral component is composed of unidentifiable plant residue with some mesquite pod. A small animal was eaten.

Table 7.8. Best determination of food components by time period.

#### Component I, Late Archaic

Sample 9: Willow catkins or foliage. Mesquite pods. Baked agave and prickly pear pads. Greens from grass and possibly from goosefoot and/or amaranth. A small animal.

#### Component II, Terminal Archaic

Sample 8: Mesquite pods and baked agave. Grass greens.

Sample 15: Willow catkins or foliage (beverage). Baked prickly pear pads. Mesquite pods. Baked agave. Greens from pigweed or goosefoot.

Sample 16: Willow catkins or foliage (beverage). Baked agave and prickly pear pads. Mesquite pods. Greens from pigweed or goosefoot. Wild squash seeds.

Sample 17: Baked agave and baked prickly pear pads. Mesquite pods. Greens from pigweed or goosefoot. Grass greens. A small animal.

Sample 18: Possibly grass.

#### Component III, Formative

Sample 1: Mesquite pods. Baked agave and prickly pear pads. Possibly goosefoot or pigweed greens. A small animal.

Sample 2: Goosefoot seed. Mesquite pods. Baked agave. Grass greens.

Sample 3: Mesquite pods. Baked agave and prickly pear pads. Greens from grass. A small animal.

Sample 4: Yucca buds or flowers. Mesquite pods. Baked prickly pear pads and agave.

Sample 5: Yucca buds or flowers. Prickly pear pads. Possibly goosefoot or pigweed greens.

Sample 6: Baked agave and prickly pear. Mesquite pods. Greens from goosefoot or pigweed.

Sample 7: Mesquite pods, baked agave, and prickly pear stems. Greens from goosefoot or pigweed.

Sample 10: Grass greens. Baked agave and prickly pear pads. A small animal.

Sample 11: Willow catkins or foliage. Baked yucca. Baked prickly pear pads.

Sample 12: Baked yucca, baked prickly pear pads. Greens from grass and possible pigweed or goosefoot. A small animal.

Sample 13: Prickly pear buds or flowers. Mesquite pods and baked prickly pear pads. Possible goosefoot or amaranth greens. Some baked yucca and a small animal.

Sample 19: Mustard greens. Mesquite pods.

Sample 20: Mormon tea (beverage). Mesquite pods. A small animal.

Sample 21: Mormon tea (beverage). Mesquite pods. A small animal.

High pollen percentages most likely result from consumption of flowers, fruits, or possibly seeds. With regard to *Salix*, it is probable that the catkins of the tree were eaten or brewed in tea. The same is true of *Ephedra*. Russell (1975) noted that the Pimas ate the baked fruits of *Yucca*. He has also noted that the Pimas ate the fruits of prickly pear. The florettes and seeds of several species of grass were consumed prehistorically. At Dust Devil Cave, Utah, *Sporobolus* was most commonly eaten (Reinhard 1985; Van Ness 1986). Chen-Am pollen was probably introduced by the consumption of flowers. Castetter has noted that *Chenopodium* flowers were eaten by the Yumas (1935). The possibility that *Quercus* was eaten has been raised; relative to this question, Russell has noted that the Pimas traded acorns from the Papagos, hulled them, and ground them into meal.

Low percentages of *Typha* pollen are present in several feces. In all likelihood, these grains were probably consumed with drinking water. However, Castetter and Bell (1951) have noted that *Typha* pollen was a favorite food for the Yuma, Maricopa, and Mojave Indians. There is a remote possibility that the *Typha* pollen is of dietary origin. Many Lovelock Cave coprolites are composed almost entirely of *Typha* pollen (Heizer and Napton 1969).

Although dietary usage has been assumed until this point, it must be noted that some of the plants noted above have medicinal value. According to Russell, the Pimas ate the raw flowers of *Yucca* as a purgative (1975). He also noted the use of *Ephedra* among the Pima as a cure for syphilis. Perhaps most important is the use of *Salix* to relieve pain (Vogel 1973). The use of "willow tea" was a widespread medicine among prehistoric peoples. Usually the tea is described as being brewed from willow bark or from the leaves of the plant. The introduction of willow pollen into willow tea could occur quite easily if catkins were fortuitously included with foliage in the brewing of tea. Willow contains salicin which is a pain reliever similar to aspirin. It is a simple acid that can be absorbed through the stomach.

Eight of the 19 human coprolites examined in this study contain pollen of plants that are potentially medicinal (Samples 9, 11, 15, 16, 20, and 21 stand out).

Seasonality is indicated by the presence of *Quercus* pollen in most coprolites and by the presence of *Salix* in some. These plants produce pollen in early spring, suggesting that the cave was used at this time.

Five coprolites contain many pollen taxa, none of which have clear dietary origin. The pollen in all of these coprolites is extremely degraded. The pollen taxa reflect the natural pollen rain one would expect in the Black Mountains. Because these represent a complete ecological spectrum, and the preservation is so poor, these grains were likely consumed with drinking water.

The Bighorn Cave coprolites are interesting palynologically because of the number of taxa present, which could allow for environmental reconstruction if botanical data were provided along with soil samples from the locality of the cave. The soil samples could be processed palynologically to serve as a comparative base for past climate reconstruction. Bighorn Cave pollen is also interesting from the perspective of paleopharmacology—the possibility that medicines are reflected in the diet is fascinating.

## COMPARISONS

Comparative analyses are available from Danger and Hogup Caves, both in Utah (Fry 1977), Lovelock Cave in Nevada (Heizer and Napton 1969), Dust Devil Cave in Utah (Reinhard 1985; Reinhard et al. 1985; Van Ness 1986), and Hinds Cave in Texas (Williams-Dean 1978; Reinhard in progress). The diets of various Southwest hunter-gatherer groups were quite variable. People at Hinds Cave depended largely on *Opuntia* seeds, *Allium* bulbs, *Opuntia* pads, *Sporobolus* seed, and walnuts. Nearly all of the 120 coprolites from that cave examined macroscopically contained bone (Williams-Dean 1978; Reinhard in progress). In Dust Devil Cave on the Colorado Plateau, Archaic people depended largely on *Opuntia* pads, *Chenopodium* seed, and *Sporobolus* seed. About 68 percent of the coprolites from Dust Devil Cave contained

bone (Reinhard 1985; Reinhard et al. 1985; Van Ness 1986). Coprolites from Danger and Hogup Caves show dependence on *Allenrolfea* seed, composite seed, and *Opuntia*. Bone was common in these coprolites as well (Fry 1977). The diet reflected in Lovelock Cave coprolites shows an adaptation to an aquatic ecosystem with consumption of fish, *Typha*, and bullrush.

The macroscopic remains from Bighorn Cave show a population that was dependent on a variety of foods. Dietary mainstays were prickly pear pads, mesquite, and coarse fiber, probably derived from *Agave* leaves. However, other plants were also important contributions to the diet, including mustard seed, chenopod seed, juniper seed, and grass seed. Whether acorns were consumed remains to be demonstrated. Bone was not very common in the coprolites (4 of 20 human coprolites contained bone); perhaps this represents a reduced consumption of meat at Bighorn Cave in comparison to other hunter-gatherer caves.

It is interesting that there is not much variation in the main diet constituents between the depositional components at Bighorn Cave (Table 7.8). Mesquite, probable *Agave*, and prickly pear were the main foods in all times. Willow, goosefoot, or pigweed greens and small animals were consumed in all components. Grass phytoliths were found in the coprolites that contain small animal bones; this suggests that these phytoliths might have been eaten by the animals and therefore grass may not have been an intentional part of the human diet. Wild squash seeds were eaten during Component II times (terminal Archaic); this dietary aspect is unique to that period. Several more constituents were present in Component III (Formative), such as goosefoot seed, mustard greens, flowers or buds of prickly pear, flowers or buds of yucca, and Mormon tea. The evident greater diversity in Component III is likely a result of having analyzed more feces from this component (15 compared to 4 for Component II and 1 for Component I). Increased sample sizes are needed for Components II and especially I before trends

among the components could have behavioral meaning.

#### PARASITE DISCUSSION

Coprolites from several Archaic, hunter-gatherer caves have been examined parasitologically in the Great Basin (Fry 1977; Heizer and Napton 1969), from the Colorado Plateau (Reinhard et al. 1985), and from the trans-Pecos of Texas (Reinhard in progress). These serve as a comparative base for the parasitological analysis of Bighorn Cave coprolites.

Parasitological examination of hunter-gatherer coprolites rarely reveals parasitism. The examination of coprolites from Lovelock Cave in Nevada revealed only free-living nematode larvae and one possible fluke infection (Heizer and Napton 1969). The examination of 100 coprolites from Dust Devil Cave in Utah revealed no parasite infections (Reinhard et al. 1985) nor did the analysis of 20 coprolites from Hinds Cave in Texas (Reinhard in progress). Only in the analyses of coprolites from Danger and Hogup Caves in Utah were parasites found. These included pinworm, thorny-headed worm, taeniid tapeworms, and head lice (Fry 1977). The lack of parasites in the Bighorn Cave coprolites is consistent with most hunter-gatherer caves. One concludes that parasitic disease was infrequent among southwestern hunter-gatherers.

Several aspects of the lives of hunter-gatherers served to limit parasitism (Reinhard 1985a). In general, hunter-gatherer populations are dilute and intergroup contact is infrequent. Usually, fecal contamination of food is limited and thus the spread of parasite diseases is limited. One main way by which hunter-gatherers became infected with parasites was by the consumption of incompletely cooked meats and insects. This was probably the source of infection for the Hogup Cave and Danger Cave populations.

Some protozoal and bacterial pathogens are transmitted by the same conditions as parasitic worms, so it is safe to conclude that disease transferred by fecal contamination in general was limited among the Bighorn

Cave inhabitants. The diversity of foods and the presence of possible medicinal plants show that the people were well adjusted to their environment.

## SUMMARY OF RECOVERED QUIDS AND FECES

*Phil R. Geib*

This chapter concerns human and other animal waste recovered from Bighorn Cave. The emphasis and key portion of this chapter was the previous detailed analysis of the micro- and macroscopic plant remains within human feces. Here I present a general summary of all consumption waste in the collections. Quids are included here because this seemed the most logical place—they are after all byproducts of eating, but are processed by the mouth and not the digestive system.

Table 7.9 gives a summary of the waste remains recovered from the test excavations of Bighorn Cave. The table rows are organized by the components and unassigned proveniences of Locus A, as well as Locus B, with all remains from that area treated as one group. The table gives three measures of abundance for each row: a count of FS numbers that contain each type of waste, an actual count of individual items (except for rodent and sheep or deer feces), and the total weight.

### Quids

The test excavations at Bighorn Cave recovered 382 quids totaling 560.6 g, which are listed by tentative identification in Table 7.10. A moderate number of the quids exhibit distinct teeth marks (Figure 7.1). Most quids are between 3 and 6 cm in length and 1.5–2.5 cm in diameter. Many are well consolidated and the individual fibers in them are extensively crushed. Nearly all of the quids appear to consist of fibers of agave or perhaps yucca. Agave seems the most likely given the other evidence found at the site for consumption of this plant, along with the abundant ethnographic documentation of agave consumption. Consumption of

yucca leaves is unlikely (rather foul tasting) and the snarled fiber masses that result from processing yucca leaves for string production are easily distinguished from chewed quids. Yucca leaves and fibers might be chewed as part of the fiber extraction process, but this should not result in the types of remains included here as quids. In agave consumption much of the cooked leaves are swallowed, but some of the more fibrous and less flavorful portions might be chewed and then expectorated.

The “Other” column includes quids of unidentifiable material, one of a chewed legume pod (perhaps mesquite), and several of probable cactus. One of the latter contains roots that appear similar to the hedgehog cactus roots observed in the plant collections (see Chapter 8). There are also quids of unknown materials, several of which were identical; other types of cactus are possible candidates. Ultimately a detailed study of the quids will provide additional information about subsistence and perhaps medicinal practices of the Bighorn Cave occupants.

The quids with meaningful provenience occur disproportionately in Component III (174 of 212 or 82% by weight). Ground moisture in some of the layers of Component I might have decomposed a portion of its quids, but differential preservation cannot be a factor with the other components, the layers of which were essentially unaffected by moisture. It could be that there was more




Figure 7.1. Quid with obvious teeth marks from Bighorn Cave.

Table 7.9. Summary of human and animal waste from Bighorn Cave (FS = count of proveniences).

Component/Locus	Quids			Animal Feces			Human Feces		
	FS	n	wt (g)	FS	n	wt (g)	FS	n	wt (g)
Locus A									
Component IV	5	19	26.3	0	0	0.0	3	8	39.9
Component III?	1	3	3.4	0	0	0.0	1	2	7.4
Component III	31	174	267.7	6	9	(46.1)	20	131	(476.2)
Component II–III	0	0	0.0	1	1	1.0	1	54	180.2
Component II?	0	0	0.0	0	0	0.0	1	13	65.8
Component II	7	13	14.3	3	5	11.3	10	51	(284.9)
Component I	2	3	1.9	2	5	1.2	10	38	(159.1)
Unassigned	19	119	178.4	1	1	7.6	11	21	276.4
Locus B	12	51	68.6	3	3	34.2	5	6	58.8
Total	77	382	560.6	16	24	(101.4)	62	324	(1548.7)

Numbers in parentheses slightly less than true total because of unavailable weights for seven analyzed feces.

Table 7.10. Preliminary identification of quids recovered from Bighorn Cave.

Component/Locus	Agave/Yucca		Other		Total	
	n	wt (g)	n	wt (g)	n	wt (g)
Locus A						
Component IV	18	24.9	1	1.4	19	26.3
Component III?	3	3.4	0	0.0	3	3.4
Component III	159	259.7	15	8.0	174	267.7
Component II	10	11.2	3	3.1	13	14.3
Component I	3	1.9	0	0.0	3	1.9
Unassigned	112	172.1	7	6.3	119	178.4
Locus B	51	68.6	0	0.0	51	68.6
Total	356	541.8	26	18.8	382	560.6

intensive exploitation of agave during the formation of the Component III layers; there is some supporting evidence for this in the plant remains for the site, because agave leaves were most abundant in Component III as well. One consideration though is that 47 percent of the quids (or 61% by weight) came from a single provenience of Component III—Layer 3b of S1E0. Thus it could be that our test units merely sampled a location of agave consumption for Component III but not the other components.

### Feces

Excavations recovered both human and animal feces from the site, with far more of the former saved than the latter. The field crew

generally collected all items that looked potentially like human feces and any carnivore feces, although the latter were few ( $n = 5$ ). Ungulate droppings were quite common to most layers, with some interfaces between strata appearing to contain concentrations of such feces. Excavators collected just a few of these specimens for the primary purpose of radiocarbon dating. Table 7.11 lists the counts and weights of animal feces by general type using the same format as Table 7.9. The two collections of rodent feces, both of which came from the burned packrat middens of Locus B (upper grotto), are bulk samples consisting of many uncounted individual pellets. Several of the sheep or deer fecal collections are also of this bulk



Table 7.11. Identification of animal feces recovered from Bighorn Cave. Bulk samples of rodent and sheep or deer feces are not individually counted.

Component/Locus	Carnivore		Sheep/Deer		Rabbit		Rodent	
	n	wt (g)	n	wt (g)	n	wt (g)	n	wt (g)
Locus A								
Component III	2	(19.8)	7	26.3	—	—	—	—
Component II–III	1	1.0	—	—	—	—	—	—
Component II	—	—	4	11.2	1	0.1	—	—
Component I	—	—	5	1.2	—	—	—	—
Unassigned	1	7.6	—	—	—	—	—	—
Locus B	1	3.9	—	—	—	—	2	30.3
Total	5	(82.3)	16	38.7	1	0.1	2	30.3

Numbers in parentheses less than true total because of unavailable weight for analyzed carnivore feces.

nature—one from Component II consisting of 10.4 g (93% of the total for this component) and three from Component III (6.7, 6.9, and 11.6 g). With an approximate average pellet weight of 0.3 g, the count of individual sheep or deer feces in each of these bulk samples can be estimated, ranging from about 22 for the smallest collection to 40 for the largest. Sediment samples collected for flotation analysis doubtless contain examples of animal feces that would provide a fairly representative collection of the types and frequencies at the site. Layer interfaces that appeared to have higher than average concentrations of sheep or deer feces would be underrepresented, because few if any float samples were collected at layer contacts.

Human feces were counted individually, even the small fragments, so the 324 specimens listed in Table 7.9 are the total number collected. These range in size from small fragments weighing just 0.4 g to one truly large specimen at 112.8 g. There were several whole or nearly whole specimens, but most were portions of various size. Some

had been squished into the underlying sediment when fresh, thereby adding on rock, sediment, charcoal, and vegetation to varying extents. Most of these extraneous adhesions were removed during the inventory phase; thus weights reflect essentially pure fecal material. Some of the human feces are heavily filled with fibers and appear to reflect a diet that consisted largely of agave. Other feces, however, seem to consist mostly of seeds and other remains. The fecal analysis reported previously provides a detailed account of the remains within 1 carnivore and 20 human feces.

Table 7.9 lists 54 feces from a single provenience as Component II–III. All of these came from the second arbitrary excavation level of the test unit at the dripline (S1E8). Many of these were point provenienced in three dimensions within the unit and there is no doubt that they originated from layers belonging to either Component II or III. More specific assignment will require direct dating, something that might eventually be quite practical and worthwhile given the relative rarity of prehistoric human feces.

## LITERATURE CITED

- Abeloe, T. N. and P. C. Hardy  
1997 Western Screech-Owls Diurnally Roosting in a Cave. *The Southwestern Naturalist* 42:349–350.
- Adam, D. P., and P. J. Mehringer, Jr.  
1975 Modern Pollen Surface Samples—An Analysis of Subsamples. *Journal of Research U.S. Geological Survey* 3:733–736.
- Adams, E. C., M. T. Stark, and D. S. Dosh  
1993 Ceramic Distribution and Exchange: Jeddito Yellow Ware and Implications for Social Complexity. *Journal of Field Archaeology* 20(1):3–21.
- Adavasio, J. M.  
1977 *Basketry Technology, a Guide to Identification and Analysis*. Aldine, Chicago.
- Aikens, C. M.  
1970 *Hogup Cave*. University of Utah Anthropological Papers No. 93, University of Utah Press, Salt Lake City.
- Aikens, C. M., and Y. T. Witherspoon  
1986 Great Basin Numic Prehistory: Linguistics, Archaeology, and Environment. In *Anthropology of the Desert West*, edited by C. J. Condie and D. D. Fowler. University of Utah Anthropological Papers No. 110, University of Utah Press, Salt Lake City.
- Andrews, P.  
1990 *Owls, Caves, and Fossils*. University of Chicago Press, Chicago.
- Andrews, P., and E. M. N. Evans  
1983 Small Mammal Bone Accumulations Produced by Mammalian Carnivore. *Paleobiology* 9:289–307.
- Anduze, R. A., D. M. Greenwald, and M. E. Walsh-Anduze  
1994 Mead to Phoenix 500kV Transmission Line Project: Data Recovery at the Overkill Site (AZ M:7:48ASM), an Archaic Site in Mohave County, Arizona. SWCA Archaeological Report No. 94-60. Ms. on file, SWCA Inc., Flagstaff.
- Barrows, D. P.  
1900 *The Ethno-Botany of the Coahuilla Indians of Southern California*. University of Chicago Press, Chicago.
- Bell, W. H., and E. F. Castetter  
1937 *The Utilization of Mesquite and Screwbean by the Aborigines in the American Southwest*. University of New Mexico Bulletin 314, Albuquerque.
- Benson, L., and R. A. Darrow  
1981 *Trees and Shrubs of the Southwestern Deserts*. 3rd ed. University of Arizona Press, Tucson.
- Berry, C. F., and M. S. Berry  
1986 Chronological and Conceptual Models of the Southwestern Archaic. In *Anthropology of the Desert West*, edited by C. J. Condie and D. D. Fowler. University of Utah Anthropological Papers No. 110, University of Utah Press, Salt Lake City.
- Bluhm, E.  
1952 Clothing and Textiles. In *Mogollon Cultural Continuity and Change, the Stratigraphic Analysis of Tularosa and Cordova Caves*, by Paul S. Martin, John B. Rinaldo, Elaine Bluhm, Hugh C. Cutler, and Roger Grange, Jr. *Fieldiana Anthropology*, Vol. 40, Chicago Natural History Museum.

- Bohrer, V. L.  
1981 Methods of Recognizing Cultural Activity from Pollen in Archaeological Sites. *The Kiva* 46:135–142.
- Bondly, G. A., and R. H. Brooks  
1973 An Archaeological Survey of the Mead-Davis-Parker 230 KV Transmission Line 2. Ms. on file, Nevada Archaeological Survey, University of Nevada, Las Vegas.
- Bonnichsen, R., and M. H. Sorg (editors)  
1979 *Bone Modification*. Center for the Study of the First Americans, Orono, Maine.
- Breternitz, D. A.  
1966 *An Appraisal of Tree-Ring Dated Pottery in the Southwest*. Anthropological Papers of the University of Arizona 10, Tucson.
- Bronk, R. C.  
1994 Analysis of Chronological Information and Radiocarbon Calibration: The Program OxCal. *Archaeological Computing Newsletter* 41:11–16.  
1995 Radiocarbon Calibration and Analysis of Stratigraphy: The OxCal Program. *Radiocarbon* 37(2):425–430.
- Brown, D. E. (editor)  
1982 Biotic Communities of the American Southwest—United States and Mexico. *Desert Plants* 4(1-4). University of Arizona, Tucson, for Boyce Thompson Southwestern Arboretum, Superior.
- Castetter, E. F.  
1935 *Uncultivated Native Plants Used As Sources of Food*. University of New Mexico Bulletin No. 266. Biological Series, Vol. 4, No. 1. Ethnobiological Studies in the American Southwest No. 1. University of New Mexico Press, Albuquerque.
- Castetter, E. F., and W. H. Bell  
1951 *Yuman Indian Agriculture*. University of New Mexico Press, Albuquerque.
- Christenson, A. L.  
2000 Petrographic Analysis of Sands, Self-tempered Clays, and Prehistoric Ceramics from the Prescott Area. In *Archaeology in West-Central Arizona, Proceedings of the 1966 Arizona Archaeological Council Prescott Conference*, edited by T. N. Motsinger, D. R. Mitchell, and J. M. McKie. Sharlot Hall Museum Press, Prescott, Arizona.
- Cole, K. L., and R. H. Webb  
1985 Late Holocene Vegetation Change in Greenwater Valley, Mojave Desert, California. *Quaternary Research* 23:227–235.
- Colton, H. S.  
1964 Principal Hopi Trails. *Plateau* 36(3):91–94.
- Colton, H. S. (editor)  
1956 *Pottery Types of the Southwest: San Juan Red Ware, Tsegi Orange Ware, Homolovi Orange Ware, Winslow Orange Ware, Awatovi Yellow Ware, Jeddito Yellow Ware, Sichomovi Red Ware*. Museum of Northern Arizona Ceramic Series No. 3C, Flagstaff.  
1958 *Pottery Types of the Southwest: Alameda Brown Ware, Tizon Brown Ware, Lower Colorado Buff Ware, Prescott Gray Ware, San Francisco Mt. Gray Ware*. Museum of Northern Arizona Ceramic Series No. 3D, Flagstaff.
- Coville, F. V.  
1892 The Panamint Indians of California. *American Anthropologist* 5(4):51–353.
- Coville, F. V.  
1893 *Botany of the Death Valley Expedition: Report on the botany of the expedition sent out in 1891 by the U. S. Department of Agriculture to make a biological survey of the region of Death Valley, California*. U.S. Dept. of Agriculture. Division of Botany. Contributions from the U.S. National Herbarium, Vol. 4. Government Printing Office, Washington, D C.
- Cosgrove, C. B.  
1947 *Caves of the Upper Gila and Hueco Areas in New Mexico and Texas*. Papers of the Peabody Museum of American Archaeology and Ethnology, Harvard University, Vol. XXIV, No. 2, Cambridge.

- Coulam, N. J., and A. R. Schroedl  
2001 Split-Twig Figurines: Late Archaic Totems in the Greater American Southwest. Ms. in review, copy obtainable from author, Bureau of Reclamation, Salt Lake City.
- Cruz-Urbe, K., and R. G. Klein  
1998 Hyrax and Hare Bones from Modern South African Eagle Roosts and the Detection of Eagle Involvement in Fossil Bone Assemblages. *Journal of Archaeological Science* 25:135–147.
- Dalley, G. F.  
1970 Artifacts of Wood. In *Hogup Cave*, by C. M. Aikens, pp. 153–186. Anthropological Paper No. 93, University of Utah Press, Salt Lake City.
- Danielson, D. R., and K. J. Reinhard  
1998 Human Dental Microwear Caused by Calcium Oxalate Phytoliths in Prehistoric Diet of the Lower Pecos Region, Texas. *American Journal of Physical Anthropology* 107:297–304.
- Davis, C. A., and G. A. Smith  
1981 *Newberry Cave*. San Bernardino County Museum Association, Redlands, California.
- Davis, O. K., and D. S. Shafer  
1992 A Holocene Climatic Record for the Sonoran Desert from Pollen Analysis of Montezuma Well, Arizona, USA. *Paleogeography, Paleoclimatology and Paleoecology* 92:107–119.
- Davis, O. K., R. H. Hevly, and R. Faust  
1985 A Comparison of Historic and Prehistoric Vegetation Change Caused by Man in Central Arizona. *AASP Contrib. Series* 16:63–75.
- Deegan, A. C.  
1994 Anasazi Fibrous Sandal Technology. *Kiva* 59:49–64.
- Dick, H. W.  
1965 *Bat Cave*. School of American Research Monograph No. 27, Santa Fe, New Mexico.
- Diveley-White, S. A.  
1987 Fauna. In *An Archaeological Evaluation of Bighorn Cave, Mohave County, Arizona*, edited by P. R. Geib and D. R. Keller, pp. 4.27–4.29. Unpublished draft ms. on file, Bureau of Land Management Office, Kingman, Arizona.
- Dobyns, H. F., and R. C. Euler  
1958 Tizon Brown Ware: A Descriptive Revision. In *Pottery Types of The Southwest*, edited by H. S. Colton. Museum of Northern Arizona Ceramic Series No. 3D. Flagstaff.  
  
1970 *Wauba Yuma's People: The Comparative Socio-political Structure of the Pai Indians of Arizona*. Prescott College Press, Prescott, Arizona.
- Dodson, P., and D. Wexlar  
1979 Taphonomic Investigation of Owl Pellets. *Paleobiology* 5:275–284.
- Dosh, S. G., M. Zyniecki, and W. S. Marmaduke  
1999 *Archaeological Investigations, U.S. Highway 93. Archaeological Sites around Kaiser Spring and along the Santa Maria River*. Northland Research Technical Report 99-07. Flagstaff.
- Driver, H. E.  
1957 Estimation of Intensity of Land Use from Ethnobiology: Applied to the Yuma Indians. *Ethnohistory* 4:174–197.
- Driver, H. E., and W. C. Massey  
1957 Comparative Studies of North American Indians. *Transactions of the American Philosophical Society* n.s. 47(2). Philadelphia.
- Drucker, P.  
1937 Culture Element Distributions. V: Southern California. *University of California Anthropological Records* 1:1–52. Berkeley.
- Emery, I.  
1966 *The Primary Structures of Fabrics, an Illustrated Classification*. The Textile Museum, Washington, D.C.
- Emslie, S. D.  
1988 Vertebrate Paleontology and Taphonomy of Caves in Grand Canyon, Arizona. *National Geographic Research* 4:128–142.

- Emslie, S. D., J. I. Mead, and L. Coats  
 1995 Split-twig Figurines in Grand Canyon, Arizona: New Discoveries and Interpretations. *Kiva* 61(2):145–173.
- Euler, R. C.  
 1958 *Walapai Culture-History*. Ph.D. dissertation, University of New Mexico, Albuquerque. University Microfilms, Ann Arbor.  
 1962 Review of *The Archaeological Excavations at Willow Beach, Arizona 1950*, by Albert H. Schroeder. *American Antiquity* 28(1):114–115.  
 1982 Ceramic Patterns of the Hakataya Tradition. In *Southwestern Ceramics, A Comparative Review*, edited by A. H. Schroeder. The Arizona Archaeologist No. 15. Phoenix.  
 1984 The Archaeology and Geology of Stanton's Cave. In *The Archaeology, Geology, and Paleobiology of Stanton's Cave, Grand Canyon National Park*, edited by R. C. Euler, pp. 7–32. Grand Canyon Natural History Association Monograph No. 6, Grand Canyon, Arizona.
- Euler, R. C., and H. F. Dobyns  
 1985 The Ethnoarchaeology of Upland Arizona Yuman Ceramics. In *Southwestern Culture History: Collected Papers in Honor of Albert H. Schroeder*, edited by C. H. Lang. The Archaeological Society of New Mexico 10. Ancient City Press, Santa Fe.
- Ferdon, E. N., Jr.  
 1946 *An excavation of Hermit's Cave, New Mexico*. School of American Research Monograph No. 10, University of New Mexico Press, Albuquerque.
- Fish, S. K.  
 1985 Prehistoric Disturbance Floras of the Lower Sonoran Desert and their Implications. *AASP Contributions Series* 16:77–88.
- Ford, R. I.  
 1983 Inter-Indian Exchange in the Southwest. In *Handbook of North American Indians, Volume 10, Southwest*, edited by A. Ortiz. Smithsonian Institution, Washington, D.C.
- Fowler, D. D., Editor  
 1973 *S. M. Wheeler, Archaeology of Etna Cave, Nevada: A Reprint*. Desert Research Institute Publications in the Social Sciences No. 7, Reno.
- Fowler, D. D., and D. B. Madsen  
 1986 Prehistory of the Southeastern Area. In *Handbook of North American Indians, Volume 11, Great Basin*, edited by W. T. d'Azevedo. Smithsonian Institution, Washington, D.C.
- Fowler, D. D., and C. S. Fowler  
 1981 *Anthropology of the Numa: John Wesley Powell's Manuscripts on the Numa Peoples of Western North America, 1868–1880*. Smithsonian Contributions to Anthropology No. 14, Washington, D.C.
- Fowler, D. D., D. B. Madsen, and E. M. Hattori  
 1973 *Prehistory of Southeastern Nevada*. Desert Research Institute Publications in the Social Sciences No. 6. Reno and Las Vegas.
- Fry, G. F.  
 1977 *Analysis of Prehistoric Coprolites from Utah*. University of Utah Anthropological Papers 97, University of Utah Press, Salt Lake City.
- Gallagher, M. V.  
 1977 *Contemporary Ethnobotany Among the Apache of the Clarkdale, Arizona Area: Coconino and Prescott National Forests*. Archaeological Report No. 14. U.S. Forest Service, Southwest Region, Albuquerque.
- Geib, P. R.  
 1986 Grinding Implement Production Near Riviera, Arizona: Archaeological Investigations at the Riviera Substation. Northern Arizona University Archaeological Report No. 760. Ms. on file, Anthropology Department, Northern Arizona University, Flagstaff.  
 1996 *Glen Canyon Revisited*. Anthropological Paper No. 119, University of Utah Press, Salt Lake City.

- Geib, P. R., and D. R. Keller (editors)  
 1987 An Archaeological Evaluation of Bighorn Cave, Mohave County, Arizona. Draft report submitted to the Bureau of Land Management. Ms. on file, Museum of Northern Arizona, Flagstaff.
- Geib, P. R., J. H. Collette, and K. Spurr  
 2001 Kaibabitsinügwü: An Archaeological Sample Survey of the Kaiparowits Plateau. Cultural Resource Series No. 25, Bureau of Land Management, Salt Lake City.
- Gray, F., R. C. Jachens, R. J. Miller, R. C. Turner, E. K. Livo, D. H. Knepper, Jr., J. Mariano, and C. L. Almquist  
 1990 *Mineral Resources of the Mount Nutt Wilderness Study Area, Mohave County, Arizona*. USGS Bulletin 1737D:1–22.
- Gray, J.  
 1965 Pollen Extraction Techniques. In *Handbook of Paleontological Techniques*, edited by B. Kummel and D. Raup, pp. 471–481. W. H. Freeman, San Francisco.
- Greenwald, D.  
 1987 Bighorn Cave: A Comparison with Other Late Archaic Lithic Assemblages. Paper presented at the Arizona Nevada Academy of Sciences Conference, April 1987, Flagstaff.
- Guernsey, S. J.  
 1931 *Explorations in Northeastern Arizona, Report on the Archaeological Fieldwork of 1920–1923*. Papers of the Peabody Museum of American Archaeology and Ethnology, Harvard University, Vol. XII, No. 1, Cambridge.
- Guernsey, S. J., and A. V. Kidder  
 1921 *Basket Maker Caves of Northeastern Arizona, Report on the Explorations, 1916–17*. Papers of the Peabody Museum of American Archaeology and Ethnology, Harvard University, Vol. VIII, No. 2, Cambridge.
- Hall, S. A.  
 1981 Deteriorated Pollen Grains and the Interpretation of Quaternary Pollen Diagrams. *Review of Paleobotany and Palynology* 32:193–206.
- Harrington, M. R.  
 n.d. *Ancient Tribes of the Boulder Dam Country*. Southwest Museum Leaflets No. 9. Southwest Museum, Highland Park, in cooperation with the Clark County Archaeological and Historical Society.
- Haury, E. W.  
 1950 *The Stratigraphy and Archaeology of Ventana Cave*. University of Arizona Press, Tucson.
- Havinga, A. J.  
 1974 An experimental investigation into the decay of pollen and spores in various soil types. In *Sporopollenin*, edited by J. Brooks et al., pp. 446–479. Academic Press, New York.
- Heizer, R. F., and L. K. Napton  
 1969 Biological and Cultural Evidence from Prehistoric Human Coprolites. *Science* 165:563–568.
- Heuett, M. L.  
 1974 *Boulder Springs: A Cerbat-Hualapai Rock Shelter in Northwestern Arizona*. Unpublished Master's thesis, Department of Anthropology, Northern Arizona University, Flagstaff.
- Hevly, R. H.  
 1981 Pollen Production, Transport and Preservation: Potentials and Limitations in Archaeological Palynology. *Journal of Ethnobiology* 1:39–54.
- Hevly, R. H., and L. E. Renner  
 1984 Pollen, Spores and Other Atmospheric Particulates, Flagstaff, Arizona. *Journal of the Arizona Academy of Science* 19:121–132.
- Hevly, R. H., P. J. Mehringer, Jr., and H. G. Yocum  
 1965 Modern Pollen Rain in the Sonoran Desert. *Journal of the Arizona Academy of Science* 3:123–135.
- Hevly, R. H., M. L. Heuett, and S. J. Olsen  
 1978 Paleoeological Reconstruction from an Upland Patayan Rock Shelter. *Journal of the Arizona-Nevada Academy of Science* 13:67–78.

- Hewitt, N. J.  
 1980 Fiber Artifacts. In *Cowboy Cave*, by Jesse D. Jennings, pp. 49–74. University of Utah Anthropological Papers No. 104, University of Utah Press, Salt Lake City.
- Hockett, B. S.  
 1991 Toward Distinguishing Human and Raptor Patterning on Leporid Bones. *American Antiquity* 56:667–679.
- Hoffman, R.  
 1988 The Contribution of Raptorial Birds to Patterning in Small Mammal Assemblages. *Paleobiology* 14:81–90.
- Holmer, R. N.  
 1978 *A Mathematical Typology for Archaic Projectile Points of the Eastern Great Basin*. Ph.D. dissertation, Department of Anthropology, University of Utah. University Microfilms, Ann Arbor.
- 1980a Chipped Stone Projectile Points. In *Cowboy Cave*, by J.D. Jennings, pp. 31–38. Anthropological Papers No. 104. University of Utah Press, Salt Lake City.
- 1980b Projectile Points. In *Sudden Shelter*, by J.D. Jennings, A.R. Schroedl, and R.N. Holmer, pp. 63–83. Anthropological Papers No. 103. University of Utah Press, Salt Lake City.
- 1986 Common Projectile Points of the Intermountain West. In *Anthropology of the Desert West. Essays in Honor of Jesse D. Jennings*, edited by C. J. Condie and D. D. Fowler, pp. 89–115. Anthropological Papers No. 110. University of Utah Press, Salt Lake City.
- Horne, P. D.  
 1985 A Review of the Evidence of Human Endoparasitism in the Pre Columbian New World Through the Study of Coprolites. *Journal of Archaeological Science* 122:299–310.
- Hough, W.  
 1914 *Culture of the Ancient Pueblos of the Upper Gila River Region, New Mexico and Arizona*. United States National Museum, Bulletin 87, Washington, D.C.
- Huckell, B. B.  
 1986 *A Ground Stone Implement Quarry on the Lower Colorado River, Northwestern Arizona*. Cultural Resource Series, Monograph No. 3. Arizona State Office, Bureau of Land Management, Phoenix.
- 1995 *Of Marshes and Maize, Preceramic Agricultural Settlements in the Cienega Valley, Southeastern Arizona*. Anthropological Papers of the University of Arizona No. 59, Tucson.
- Hughes, R. E., and J. A. Bennyhoff  
 1986 Early Trade. In *Handbook of North American Indians, Volume 11, Great Basin*, edited by W. L. d'Azevedo. Smithsonian Institution, Washington, D.C.
- Hurley, W. M.  
 1979 *Prehistoric Cordage, Identifications of Impressions on Pottery*. Taraxacum Inc., Washington, D.C.
- Hurst, C. T.  
 1944 1943 Excavation in Cave II, Tabeguache Canyon, Montrose County, Colorado. *Southwestern Lore* 10:2–14.
- 1947 Excavations of Dolores Cave – 1946. *Southwestern Lore* 13:8–17.
- 1948 The Cottonwood Expedition, 1947 – A Cave and a Pueblo Site. *Southwestern Lore* 14:4–19.
- Jacobs, B. F.  
 1984 Identification of Southwestern Pine Species Applied to a Pollen Record from Hay Lake, Arizona. Sixth International Palynological Conference (IPC), Calgary, Alberta, Canada, *Abstracts* 71.
- Janetski, J. C.  
 1980 Wood and Reed Artifacts. In *Cowboy Cave*, by Jesse D. Jennings, pp. 75–96. University of Utah Anthropological Papers No. 104, University of Utah Press, Salt Lake City.

- Jass, C. N.  
2000 *Mid- to Late-Holocene Mammalian Fauna and Site Taphonomy of Bighorn Cave, Black Mountains, Arizona*. Unpublished Masters thesis, Northern Arizona University, Flagstaff.
- Jass, C. N.  
n.d. Mid- to Late-Holocene Mammals from Bighorn Cave, Arizona. Submitted to *Southwestern Naturalist*.
- Jennings, J. D.  
1957 *Danger Cave*, edited by R. B. Woodbury. Memoirs of the Society for American Archaeology, No. 14, published jointly by University of Utah Press and Society for American Archaeology.  
1980 *Cowboy Cave*. University of Utah Anthropological Papers No. 104, University of Utah Press, Salt Lake City.
- Jennings, J. D., and E. Norbeck  
1955 Great Basin Prehistory: A Review. *American Antiquity* 21:1–11.
- Keller, D. R.  
1983 Mead-Phoenix 500 KV DC Transmission Corridor Sensitivity Studies: Archaeology and History. Ms. on file, Museum of Northern Arizona, Flagstaff.  
1985 Archaeological Survey of the Mead to Phoenix 500 kV Direct Current Transmission Line Preferred Alternative. Ms. on file, Museum of Northern Arizona, Flagstaff.  
1992 An Archaeological Survey of a Proposed 3.7 Mile Fenceline on Prospect Ridge, Hualapai Indian Reservation. Ms. on file, Hualapai Wildlife Management Department, Peach Springs (Cultural Resources Report No. 3).  
1993 San Francisco Mountain Gray Ware; Prescott Gray Ware; Tizon Brown Ware; Lower Colorado Buff Ware. Contributions in *Across the Colorado Plateau: Anthropological Studies for the Transwestern Pipeline Expansion Project*, Vol. XVI, *Interpretation of Ceramic Artifacts*, by Barbara J. Mills, Christine E. Goetze, and Maria Nieves Zedeno, pp. 62–69, 80–84. Office of Contract Archeology and Maxwell Museum of Anthropology, Albuquerque.
- Kelly, R. E.  
1966 Split-Twig Figurines from Sycamore Canyon, Central Arizona. *Plateau* 38(3):65–67.
- Kelly, I. T., and C. S. Fowler  
1986 The Southern Paiute. In *Handbook of North American Indians, Volume 11, Great Basin*, edited by W. L. d'Azevedo. Smithsonian Institution, Washington, D.C.
- Kelso, G. K.  
1976 *Absolute Pollen Frequencies Applied to the Interpretation of Human Activities in Northern Arizona*. Ph.D. dissertation, University of Arizona, Tucson.
- Kidder, A. V., and S. J. Guernsey  
1919 *Archaeological Explorations in Northeastern Arizona*. Smithsonian Institution, Bureau of American Ethnology Bulletin 65, Washington, D.C.
- Klein, R. G., and K. Cruz-Urbe  
1984 *The Analysis of Animal Bones from Archeological Sites*. University of Chicago Press, Chicago.
- Kroeber, A. L.  
1902 Preliminary Sketch of the Mohave Indians. *American Anthropologist* 4:276–285.  
1925 *Handbook of the Indians of California*. Smithsonian Institution, Bureau of American Ethnology Bulletin 78, Washington, D.C.
- Kroeber, A. L., Editor  
1935 *Walapai Ethnography*. Memoirs of the American Anthropological Association No. 42. Menasha.
- Kusmer, K. D.  
1990 Taphonomy of Owl Pellet Deposition. *Journal of Paleontology* 64:629–637.
- Lehr, J. H.  
1978 *A Catalogue of the Flora of Arizona*. Desert Botanical Garden, Phoenix, Arizona.



- Lindsay, A. J., J. R. Ambler, M. A. Stein, and P. M. Hobler  
 1968 *Survey and Excavations North and East of Navajo Mountain, Utah, 1959–1962*. Museum of Northern Arizona Bulletin No. 45, Glen Canyon Series No. 8, Flagstaff.
- Loud, L. L., and M. R. Harrington  
 1929 *Lovelock Cave*. University of California Publications in American Archaeology and Ethnology, Vol. 25, No. 1. University of California Press, Berkeley.
- Lyneis, M. M.  
 1988 Tizon Brown Ware and the Problems Raised by Paddle-and-Anvil Pottery in the Mojave Desert. *Journal of California and Great Basin Anthropology* 10(2):146–155.  
 1992 *The Main Ridge Community at Lost City, Virgin Anasazi Architecture, Ceramics, and Burials*. University of Utah Anthropological Papers No. 117, University of Utah Press, Salt Lake City.
- MacMahon, R. H.  
 1985 *Deserts*. Knopf, New York.
- MacMahon, R. H., and F. H. Wagner  
 1985 The Mojave, Sonoran and Chihuahuan Deserts of North America. In *Ecosystems of the World 12A: Hot Deserts and Arid Shrublands*, edited by M. Evenari, I. Noy-Meir, and D. W. Goodall, pp. 105–202. Elsevier Publishing, Amsterdam.
- Magers, P. C.  
 1986 Weaving at Antelope House. In *Archaeological Investigations at Antelope House*, Don P. Morris. National Park Service, Washington D.C.
- Martin, P. S.  
 1963 *The Last 10,000 Years: A Fossil Pollen Record of the American Southwest*. University of Arizona Press, Tucson.
- Martin, P. S., and F. W. Sharrock  
 1964 Pollen Analysis of Prehistoric Human Feces; New Approach to Ethnobotany. *American Antiquity* 30(2):168–180.
- Matson, R. G.  
 1971 *Adaptation and Environment in the Cerbat Mountains, Arizona*. Unpublished Ph.D. dissertation, Department of Anthropology, University of California, Davis.  
 1974 The Determination of Archaeological Structure: An Example from the Cerbat Mountains, Arizona. *Plateau* 47:26–40.  
 1991 *The Origins of Southwestern Agriculture*. University of Arizona Press, Tucson.
- McClellan, C., D. A. Phillips, Jr., and M. Belshaw  
 1980 *The Archeology of Lake Mead National Recreation Area: An Assessment*. Western Archeology Center Publications in Anthropology No. 9. Tucson, Arizona.
- McGuire, T. R.  
 1983 Walapai. In *Handbook of North American Indians, Vol. 10, Southwest*, edited by A. Ortiz. Smithsonian Institution, Washington, D.C.
- McKeel, S.  
 1935 Plant Foods and Preparation. In *Walapai Ethnography*, edited by A. L. Kroeber. Memoirs of the American Anthropological Association No. 42, Menasha.
- Mehring, P. J., Jr.  
 1967 Pollen Analysis of the Tule Springs Site Area, Nevada. In *Pleistocene Studies in Southern Nevada*, edited by H. M. Wormington and D. Ellis. Nevada State Museum Anthropological Papers 13:19–200.
- Mehring, P. J., Jr.  
 1977 Great Basin Late Quaternary Environments and Chronology. In *Models and Great Basin Prehistory: A Symposium*, edited by D. D. Fowler. Desert Research Institute Publications in the Social Sciences 12:113–167.

- Mehring, P. J., Jr., and C. N. Warren.  
1976 *Marsh, Dune and Archaeological Chronology, Ash Meadows, Amargosa Desert, Nevada*. Nevada Archaeological Survey Research Paper: Holocene Environmental Change in the Great Basin 6:121–150.
- Morss, N.  
1931 *The Ancient Culture of the Fremont River in Utah*. Papers of the Peabody Museum of American Archaeology and Ethnology, Harvard University, Vol. XII, No. 3, Cambridge.
- Muto, G. R.  
1971 *A Technological Analysis of the Early Stages in the Manufacture of Lithic Artifacts*. Masters thesis, Department of Anthropology, Idaho State University, Pocatello.
- Nations, D., and E. Stump  
1996 *Geology of Arizona*. 2nd ed. Kendall/Hunt, Dubuque, Iowa.
- Olson, L. D.  
1973 *Outdoor Survival Skills*. Brigham Young University Press, Provo, Utah.
- Olson, A. P.  
1966 Split-Twig Figurines from NA5607, Northern Arizona. *Plateau* 38(3):55–64.
- Pendleton, L. S. A.  
1985 Chapter 14. Material Culture: Artifacts of Stone. In *The Archaeology of Hidden Cave, Nevada*, edited by D. H. Thomas, pp. 183–218. Anthropological Papers of the American Museum of Natural History, Vol. 61: Part 1. New York.
- Phagan, C. J. and T. H. Hruby  
1984 Reductive Technologies Manual: Preliminary Analysis Systems and Procedures. Dolores Archaeological Program Technical Reports DAP-150. Ms. on file, Bureau of Reclamation, Upper Colorado Region, Salt Lake City.
- Pilles, P. J., Jr.  
1981 A Review of Yavapai Archaeology. In *The Protohistoric Period in the North American Southwest, AD 1450–1700*, edited by D. R. Wilcox and W. B. Masse. Arizona State University Anthropological Research Papers No. 24, Tempe.
- Rea, A. M.  
1997 *At the Desert's Green Edge: An Ethnohistory of the Gila River Pima*. University of Arizona Press, Tucson.
- Reinhard, K. J.  
1985a *Recovery of Helminths from Prehistoric Feces: The Cultural Ecology of Ancient Parasitism*. M.A. thesis, Department of Biology, Northern Arizona University, Flagstaff.  
1985b *Strongyloides Stercoralis* in the Prehistoric Southwest. In *Health and Disease in the Prehistoric Southwest*, edited by C. F. Merbs and R. J. Miller, pp. 234–242. Arizona State University Anthropological Research Papers No. 34, Tempe.  
1985c Parasitism at Antelope House: A Puebloan Village in Canyon DeChelly, Arizona. In *Health and Disease in the Prehistoric Southwest*, edited by C. F. Merbs and R. J. Miller, pp. 220–223. Arizona State University Anthropological Research Papers No. 34, Tempe.  
n.d. The Cultural Ecology of Prehistoric Parasitism on the Colorado Plateau As Evidenced by Coprology. *American Journal of Physical Anthropology*.  
1988 Cultural Ecology of Prehistoric Parasitism on the Colorado Plateau As Evidenced by Coprology. *Am J Phys Anthropol* 77:355–366.
- Reinhard, K. J., and V. M. Bryant, Jr.  
1992 Coprolite Analysis: A Biological Perspective on Archaeology. In *Advances in Archaeological Method and Theory* 4, edited by M. B. Schiffer. University of Arizona Press, Tucson.
- Reinhard, K. J., J. R. Ambler, and M. McGuffie  
1985 Diet and Parasitism at Dust Devil Cave. *American Antiquity* 50:819–824.
- Reinhard, K. J., U. E. Confalonieri, B. Hermann, L. F. Ferreria, and A. D. G. de Araujo  
1986 Aspects of Paleoparasitological Technique: Recovery of Parasite Remains from Coprolites and Latrines. *Homo*.

- Reinhard, K. J., R. H. Hevly, and G. A. Anderson  
1987 Helminth Remains from Prehistoric Indian Coprolites on the Colorado Plateau. *Journal of Parasitology*. In press.
- Reinhard, K. J., R. H. Hevly, S. Clay-Poole, and C. H. Halbirt  
1985 Pollen Analysis of Dust Devil Cave Feces. In *Recovery of Helminths from Prehistoric Feces: The Cultural Ecology of Prehistoric Parasitism*, by K. J. Reinhard. M.A. Thesis, Department of Biology, Northern Arizona University, Flagstaff.
- Reynolds, S. J.  
1988 *Geologic Map of Arizona*. Arizona Geological Survey, in cooperation with U.S. Geological Survey.
- Riley, C. L.  
1982 *The Frontier People: The Greater Southwest in the Protohistoric Period*. Center for Archaeological Investigations Occasional Paper No. 1, Southern Illinois University, Carbondale.
- Russell, F.  
1975 *The Pima Indians*. University of Arizona Press, Tucson.
- Schaafsma, P.  
1980 *Indian Rock Art of the Southwest*. School of American Research, Santa Fe, and University of New Mexico Press, Albuquerque.
- Schiffer, M. B.  
1986 Radiocarbon Dates and the "Old Wood" Problem: The Case of the Hohokam Chronology. *Journal of Archaeological Science* 13:13–50.
- Schilz, A. J.  
1982 Archaeological Investigations in the Hualapai Valley, Arizona: The Pataya Gas Storage Facility. Ms. on file, Bureau of Land Management, Phoenix, and WESTEC Services, Inc., San Diego.
- Schoenwetter, J., and J. A. Doerschlag  
1971 Surficial Pollen Records from Central Arizona, I: Sonoran Desert Scrub. *Journal of the Arizona Academy of Science* 6:216–221.
- Schoenwetter, J., and F. W. Eddy  
1964 *Alluvial and Palynological Reconstructions of Environments in the Navajo Reservoir District*. Museum of New Mexico Papers in Anthropology 13:155.
- Schroeder, A. H.  
1958 Lower Colorado Buff Ware: A Descriptive Revision. In *Pottery Types of The Southwest*, edited by H. S. Colton. Museum of Northern Arizona Ceramic Series No. 3D. Flagstaff.  
1961 *The Archeological Excavations at Willow Beach, Arizona 1950*. University of Utah Anthropological Papers No. 50, University of Utah Press, Salt Lake City.
- Schroedl, A. R.  
1976 *The Archaic of the northern Colorado Plateau*. Ph. D. dissertation, University of Utah. University Microfilms International, Ann Arbor
- Schroedl, A. R., and N. J. Coulam  
1994 Cowboy Cave Revisited. *Utah Archaeology* 7(1):1–34.
- Schwartz, D. W., A. L. Lange, and R. DeSaussure  
1958 Split Twig Figurines in the Grand Canyon. *American Antiquity* 23:264–274.
- Shackley, M. S.  
1996 Elko or San Pedro? A Quantitative Analysis of Late Archaic Projectile Points from White Tanks, Yuma County, Arizona. *Kiva* 61:413–432.
- Shott, M. J.  
1997 Stones and Shafts Redux: The Metric Discrimination of Chipped-Stone Dart and Arrow Points. *American Antiquity* 62:86–101.
- Smith, W.  
1971 *Painted Ceramics of the Western Mound at Awatovi*. Papers of the Peabody Museum of Archaeology and Ethnology 38, Cambridge.

- Solomon, A. M., and H. D. Hayes  
 1972 Desert Pollen Production, I: Qualitative Influence of Moisture. *Journal of the Arizona Academy of Science* 7:52–74.
- Solomon, A. M., T. J. Blasing, and J. A. Solomon  
 1982 Interpretation of Floodplain Pollen in Alluvial Sediments from an Arid Region. *Quaternary Research* 18:52–71.
- Spicer, E. H.  
 1962 *Cycles of Conquest, the Impact of Spain, Mexico, and the United States on the Indians of the Southwest, 1533–1960*. University of Arizona Press, Tucson.
- Spier, L.  
 1928 *Havasupai Ethnography*. Anthropological Papers of the American Museum of Natural History 29(3), New York.  
 1955 *Mohave Culture Items*. Northern Arizona Society of Science and Art, Flagstaff.
- Stewart, K. M.  
 1983 Mohave. In *Handbook of North American Indians, Volume 10, Southwest*, edited by A. Ortiz. Smithsonian Institution, Washington, D.C.
- Stone, C. L.  
 1981 Economy and Warfare Along the Lower Colorado River. In *The Protohistoric Period in the North American Southwest, AD 1450–1700*, edited by D. R. Wilcox and W. B. Masse. Arizona State University Anthropological Research Papers No. 24, Tempe.  
 1986 *Deceptive Desolation: Prehistory of the Sonoran Desert in West Central Arizona*. Cultural Resource Series No. 1, Bureau of Land Management, Arizona, Phoenix.  
 1987 *People of the Desert, Canyons and Pines: Prehistory of the Patayan Country in West Central Arizona*. Cultural Resource Series No. 5, Bureau of Land Management, Arizona, Phoenix.  
 1991 *The Linear Oasis: Managing Cultural Resources Along the Lower Colorado River*. Cultural Resource Series No. 6, Bureau of Land Management, Arizona, Phoenix.
- Swarthout, J.  
 1981 An Archaeological Overview for the Lower Colorado River Valley, Arizona, Nevada, and California, Reach 2, Grand Wash Cliffs to Davis Dam. Ms. on file, Lake Mead National Recreation Area, Boulder City, and Museum of Northern Arizona, Flagstaff.  
 1986 *Strike Up the Band: Trickle-Down Hunter-Gatherer Strategies in the Cerbat Region of Arizona*. Unpublished Ph.D. dissertation, Department of Anthropology, Arizona State University, Tempe.
- Swarthout, J., and C. E. Drover  
 1981 An Archaeological Overview for the Lower Colorado River Valley, Arizona, Nevada, and California, Reach 3, Davis Dam to the International Border. Ms. on file, Lake Mead National Recreation Area, Boulder City, and Museum of Northern Arizona, Flagstaff.
- Swift, S. L., J. I. Mead, and C. N. Jass  
 1999 Holocene Squamates from an Owl Roost Deposit, Northwestern Arizona. *Journal of Vertebrate Paleontology* 19 (Supplement to No. 3):80A.
- Thomas, D. H. and S. L. Bierwirth  
 1983 Material Culture of Gatecliff Shelter: Projectile Points. In *The Archaeology of Monitor Valley. 2. Gatecliff Shelter*. Anthropology Papers Vol. 59: Part 1. The American Museum of Natural History, New York.
- Thompson, R. S.  
 1990 Vegetation and Climate in the Great Basin. In *Packrat Middens: The Last 40,000 Years of Biotic Change*, edited by J. L. Betancourt, T. R. Van Devender, and P. S. Martin, pp. 200–223. University of Arizona Press, Tucson.
- Thorson, J.P.  
 1971 *Igneous Petrology of the Oatman District, Mohave County Arizona*. Ph.D. dissertation, University of California at Santa Barbara.

- Turner, R. M.  
1982 Mohave Desertscrub. In *Biotic Communities of the American Southwest - United States and Mexico*, edited by D. E. Brown, pp. 157–168. *Desert Plants* 4(1-4).
- Turner, R. M., J. E. Bowers, and T. L. Burgess  
1995 *Sonoran Desert Plants: An Ecological Atlas*. University of Arizona Press, Tucson.
- Van Devender, T. R., and W. G. Spaulding  
1979 Development of Vegetation and Climate in the Southwestern United States. *Science* 204:701–710.
- Van Ness, M. A.  
1986 *Desha Complex Macrobotanical Fecal Remains : An Archaic Diet in the American Southwest*. Masters thesis, Department of Anthropology, Northern Arizona University, Flagstaff.
- Vogel, V. J.  
1970 *American Indian Medicine*. University of Oklahoma Press, Norman.
- Walling, B. A., R. A. Thompson, G. F. Dalley, D. G. Weder, K. M. Heath, and L. W. Lindsay  
1986 *Excavations at Quail Creek*. Cultural Resource Series No. 20, Bureau of Land Management, Arizona, Phoenix.
- Warren, C. N.  
1984 The Desert Region. In *California Archaeology*, edited by M. J. Moratto. Academic Press, New York.
- Warren, C. N., and R. H. Crabtree  
1986 Prehistory of the Southwestern Area. In *Handbook of North American Indians, Volume 11, Great Basin*, edited by W. L. d'Azevedo. Smithsonian Institution, Washington, D.C.
- Waters, M. R.  
1982 The Lowland Patayan Ceramic Tradition. In *Hohokam and Patayan, Prehistory of Southwestern Arizona*, edited by R. H. McGuire and Michael B. Schiffer. Academic Press, New York.
- Weaver, D. R.  
1991 An Archaeological Survey of Three Dam Site Parcels Within the Kaibab National Forest, Williams and Chalendar Ranger Districts, Coconino County, Arizona. Ms. on file, Kaibab National Forest, Williams, and Plateau Mountain Desert Research, Flagstaff.
- Weber, S. A., and P. D. Seaman, Editors  
1985 *Havasupai Habitat, A. F. Whiting's Ethnography of a Traditional Indian Culture*. University of Arizona Press, Tucson.
- Wendrich, W.  
1991 *Who Is Afraid of Basketry*. Centre of Non-Western Studies, Leiden University. Leiden, The Netherlands.
- Wheeler, S. M.  
1942 *Archaeology of Etna Cave, Lincoln County, Nevada*. Nevada State Park Commission, Carson City.
- Williams D. G.  
1978 *Ethnobotany and Cultural Ecology of Prehistoric Man in Southwest Texas*. Anthropological Research Laboratory, Texas A&M University, College Station.
- Worthington, A.  
1981 An Archaeological Survey of Selected 40 Acre Quadrats Within Hualapai Timber Sales II and III. Ms. on file, Anthropological Laboratory, Northern Arizona University.
- Wright, B. A.  
1954 *Excavation of Catclaw Cave, Lower Colorado River*. Unpublished M.A. thesis, Department of Anthropology, University of Arizona, Tucson.
- Wright, T. E., and D. E. Purcell  
2000 An Archaeological Transect Across the Western Prescott Periphery: Prehistoric Site Distribution Based on Combined Survey Results. In *Archaeology in West-Central Arizona, Proceedings of the 1966 Arizona Archaeological Council Prescott Conference*, edited by T. N. Motsinger, D. R. Mitchell, and J. M. McKie. Sharlot Hall Museum Press, Prescott.